Type B aortic dissection in a chronic haemodialysis patient

Miguel Bigotte Vieira,¹ Tiago Ferreira,² Patricia Cotovio,³ Augusto Ministro²

¹Nephrology Department, Hospital de Santa Maria—Centro Hospitalar Lisboa Norte, Lisbon, Portugal

²Vascular Surgery Department, Hospital de Santa Maria—Centro Hospitalar Lisboa Norte, Lisbon, Portugal

³Nephrology Department, Hospital Curry Cabral—Centro Hospitalar Lisboa Central, Lisbon, Portugal

Correspondence to
Dr Miguel Bigotte Vieira,
mbigottevieira@gmail.com

Accepted 1 November 2015

DESCRIPTION

A 61-year-old African-American patient presented to the emergency room with severe back pain. The patient had a history of end-stage renal disease on haemodialysis for 6 years, chronic obstructive lung disease and uncontrolled hypertension. CT scan revealed type B aortic dissection extending from the left subclavian artery to the right renal artery

(figures 1–3). There were no signs or symptoms of malperfusion. The patient was medically managed; blood pressure control required four different classes of antihypertensive drugs. It should be noted that the patient had consistently refused anti-hypertensive treatment previously. At 5 months of follow-up, the patient was asymptomatic and the dissection was stable.

Figure 1 CT scan—aortic dissection extending from left subclavian artery to right renal artery. (A) Coronal plane. (B) Sagittal plane.

Figure 2 CT three-dimensional reconstruction with true and false lumens (arrows) extending from left subclavian artery to right renal artery. (A) Posterior incidence. (B) Right lateral incidence. (C) Anterior incidence.

To cite: Bigotte Vieira M, Ferreira T, Cotovio P, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-213376

Figure 3 CT three-dimensional reconstruction with true and false lumens (arrows) extending from left subclavian artery to right renal artery. Left lateral incidence. (A–D) Four transverse plane sections.

Learning points

- ▶ Aortic dissection, which is more common in men and older people, has many predisposing factors including hypertension (most common), male gender, older age, vasculitis, disorders of collagen, bicuspid aortic valve, aortic coarctation and cocaine use.
- ▶ According to the Stanford classification, which has treatment and prognostic implications, aortic dissection can be categorised as type A, when the ascending aorta is involved, or as type B, when aortic dissection occurs distally to the left subclavian artery.²
- ▶ Aortic dissection diagnosis in haemodialysis patients requires a high index of suspicion as clinical presentation may be suggestive of many other conditions and only a small number of cases have been reported in this population.³

Aortic dissection is characterised by separation of the layers within the aortic wall. The incidence of acute aortic dissection is estimated to range from 3.5 to 14 per 100 000 person-years while in-hospital mortality is 27.4%.^{1 2} Although the clinical presentation might include aortic regurgitation, cardiac tamponade and end-organ ischaemia, the single most common symptom is sudden onset of severe chest pain.² Notably, the most common predisposing factor is a history of hypertension, being present in more than 70% of patients and being more common in type B dissection. Although patients with type B dissection present decreased mortality when compared to type A dissection patients, they tend to be older, have higher rates of atherosclerosis and are more likely to present with back pain rather than anterior chest pain.² Treatment options include medical therapy comprising β -blockers and surgical or endovascular intervention.

Acknowledgements The authors thank the CT radiographer, Rui Araújo, for three-dimensional image processing.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Patel AY, Eagle KA, Vaishnava P. Acute type B aortic dissection: insights from the International Registry of Acute Aortic Dissection. *Ann Cardiothorac Surg* 2014;3:368–74.
- 2 Hagan PG, Nienaber CA, Isselbacher EM, *et al*. The International Registry of Acute Aortic Dissection (IRAD): new insights into an old disease. *JAMA* 2000;283:897–903.
- 3 Tong Y, Wang Q, Hou Z. Aortic Dissection and Hemodialysis: A Case Report. *Dial Transplant* 2007;36:543–51.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow