SAPHO syndrome with mandibular manifestation

Akira Baba,¹ Hiroya Ojiri,² Shinichi Takahashi,³ Akira Katakura⁴

¹Department of Radiology, Tokyo Dental College, Ichikawa General Hospital, Chiba, Japan

²Department of Radiology, The Jikei University School of Medicine and University Hospital, Tokyo, Japan

³Department of Dermatology, Tokyo Dental College, Ichikawa General Hospital, Chiba, Japan

⁴Department of Oral Pathobiological Science and Surgery, Tokyo Dental College, Tokyo, Japan

Correspondence to

Dr Akira Baba,
akirababa@jikei.ac.jp

Accepted 12 December 2015

DESCRIPTION

A 47-year-old man was referred to our hospital due to symptoms of trismus and exacerbation of palmo-plantar pustulosis. He had been presenting, for 20 years, with refractory and recurrent mandibular osteomyelitis accompanied by palmoplantar pustulosis and anterior chest pain—symptomatic treatment and surgery had been given at each

Figure 1 MRI (short tau inversion recovery) showed high signal intensity in the right ramus of the mandible, with perilesional soft tissue swelling.

presentation. When the mandibular lesion got worse this time, he also had anterior chest pain. He had experienced similar episodes before this event. MRI (STIR: short tau inversion recovery) showed high signal intensity in the right ramus of the mandible, with perilesional soft tissue swelling ([figure 1](#)). These findings indicated active mandibular osteomyelitis. SAPHO syndrome was suggested by the clinical course and history. Bone scintigraphy using ^{99m}Tc-methylenediphosphonate (MDP) was performed and increased tracer uptake was seen in the right ramus of the mandible, greater wing of sphenoid bone, sternum and the sternocostoclavicular joints ([figure 2](#)). These characteristic clinical and radiological findings were compatible with the diagnosis of SAPHO syndrome involving the mandibular region. The term SAPHO syndrome was coined in 1987 to describe an association of characteristic bone, joint and skin lesions.¹ ‘SAPHO’ is an acronym for synovitis, acne, palmoplantar pustulosis, hyperostosis and osteitis. Its aetiology remains unknown.^{1 2} The sternocostoclavicular, sternoclavicular, costoclavicular and manubriosternal joints are affected.³ Bone scintigraphy using ^{99m}Tc-MDP is known to be important for diagnosis of this disease. SAPHO syndrome in the mandibular region is clinically rare and has been reported to be 10% of this entity.¹ SAPHO syndrome in the mandibular region is seen as recurrent mandibular osteomyelitis, mainly involving the body of the mandible.² In patients with clinical and radiological evidence of SAPHO syndrome, when recurrent mandibular

Figure 2 Bone scintigraphy using ^{99m}Tc-methylenediphosphonate revealed increased tracer uptake in the right ramus of the mandible, greater wing of sphenoid bone, sternum and the sternocostoclavicular joints.

To cite: Baba A, Ojiri H, Takahashi S, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-213401

osteomyelitis is also present, it should be considered as a part of SAPHO syndrome.

Learning points

- ▶ SAPHO syndrome involving the mandible is clinically rare and has been reported to be 10% of this entity.
- ▶ SAPHO syndrome in the mandibular region is seen as recurrent mandibular osteomyelitis, mainly involving the body of the mandible.
- ▶ In patients with evidence of SAPHO syndrome, when recurrent mandibular osteomyelitis is also present, it should be considered as a part of SAPHO syndrome.

Contributors AB drafted the paper, and collected data and reviewed the text. All the authors diagnosed the disease, read and approved the final manuscript. ST and AK treated the patient and helped AB draft the paper. HO made the final revision.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Chamot AM, Benhamou CL, Kahn MF *et al.* [Acne-pustulosis-hyperostosis-osteitis syndrome. Results of a national survey. 85 cases]. *Rev Rhum Mal Osteoartic* 1987;54:187–96.
- 2 Zemann W, Pau M, Feichtinger M *et al.* SAPHO syndrome with affection of the mandible: diagnosis, treatment, and review of literature. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;111:190–5.
- 3 Kundu BK, Naik AK, Bhargava S *et al.* Diagnosing the SAPHO syndrome: a report of three cases and review of literature. *Clin Rheumatol* 2013;32:1237–43.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow