

Excision of a distended chronic non-functional large ileoanal pouch

Phillip Correia Copley,¹ Robert Tyler,² Atif Alvi²

¹Department of Trauma & Orthopaedics, Addenbrooke's Hospital, Cambridge, UK
²Department of Colorectal Surgery, Hinchingbrooke Hospital, Huntingdon, UK

Correspondence to
Dr Phillip Correia Copley,
phillip.copley@nhs.net

Accepted 23 December 2015

DESCRIPTION

Restorative proctocolectomy with ileal pouch-anal anastomosis is the surgical treatment of choice for patients with refractory ulcerative colitis. Long-term complications of this procedure include pouchitis, metabolic complications, strictures, fistulae, pouch failure and pouch neoplasia.^{1 2} One series of 1911 patients with ileal pouch-anal anastomosis found a 3.5% failure rate at 5.5 years with causes including: pelvic sepsis, high stool volume, incontinence and disease remission.³ We present a rare case of massive abdominal distension due to non-functioning of the pouch necessitating complete excision of the pouch with end ileostomy to alleviate intractable symptoms.

A 73-year-old woman with a history of ulcerative colitis and colectomy with ileoanal pouch formation almost 30 years prior was experiencing chronic abdominal distension, diarrhoea and malnutrition. She had lost 10 kg over 1 year and her body mass index had dropped to 15 kg/m². Examination revealed a hugely distended abdomen with visible peristalsis and tympanic resonance on percussion. Imaging demonstrated a grossly distended pouch and proximal bowel dilation (figures 1 and 2). Excision of the non-functional pouch was performed via a midline laparotomy. Findings were an abnormally dilated non-functional pouch with an approximate size of 44×22 cm (figure 3), and grossly dilated small bowel up to the proximal jejunum with no point of obstruction. The pouch was mobilised up to the pelvic floor and completely excised at the anorectal junction. A defunctioning right iliac fossa ileostomy was fashioned. Postoperatively, a prolonged stay on intensive care



Figure 2 Sagittal view of CT scan demonstrating the extent of this massively distended ileoanal pouch.




Figure 3 Intraoperative photograph demonstrating chronically distended, large, non-functional ileoanal pouch with dilation of the proximal small bowel.



Figure 1 Abdominal radiograph demonstrating hugely dilated ileoanal pouch with proximal small bowel dilation.

was necessary due to difficulty weaning from mechanical ventilation. The patient went on to make a good recovery.

Learning points

- ▶ Chronic distension of an ileoanal pouch can lead to malabsorption syndrome.
- ▶ Complete excision of the pouch with creation of a defunctioning end ileostomy may be curative in such patients.

Contributors PCC wrote the article, and RT and AA reviewed the article.

Competing interests None declared.



To cite: Copley PC, Tyler R, Alvi A. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-213662

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Shen B. Diagnosis and management of postoperative ileal pouch disorders. *Clin Colon Rectal Surg* 2010;23:259–68.
- 2 Gorgun E, Remzi F. Complications of ileoanal pouches. *Clin Colon Rectal Surg* 2004;17:43–55.
- 3 Delaney CP, Remzi FH, Gramlich T, *et al*. Equivalent function, quality of life and pouch survival rates after ileal pouch-anal anastomosis for indeterminate and ulcerative colitis. *Ann Surg* 2002;236:43–8.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow