Spontaneous subcutaneous tissue haematoma associated with warfarin

Nuno Cercas Pinheiro,¹ Ana Lopes,¹ Ana Camões,² Micaela Seemann Monteiro²

¹Department of Medicine I, Hospital Egas Moniz, CHLO, Lisbon, Portugal

²Emergency Department H. São Francisco Xavier, CHLO, Lisbon, Portugal

Correspondence to

Dr Nuno Cercas Pinheiro, ncercaspinheiro@gmail.com

Accepted 24 March 2016

DESCRIPTION

A 75-year-old man with a history of non-valvular atrial fibrillation anticoagulated on warfarin for the past 5 years without any prior complications presented to the emergency department with a cutaneous haemorrhagic lesion on his right lower limb, developed within the previous 24 h. There was no history of local skin lesion or trauma, and no infection or erroneous warfarin overdose.

Physical examination highlighted voluminous haematic phlyctena in the anterior, lateral surface of the right leg, surrounded by a large haematoma (figure 1A, B). Laboratory results revealed a supra therapeutic international normalised ratio (INR) (5.9), with no evidence of anaemia, nor of hepatic or renal abnormalities, and no signs of infection. A soft tissue ultrasound was performed revealing a large and heterogeneous haematoma of the local subcutaneous tissue, without muscular involvement, measuring 15×5 cm, compatible with the diagnosis of acute/subacute haematoma.

Drainage of the haematic phlyctena was then performed, as well as correction of INR with fitomenadione and human prothrombin complex. Self reabsorption of the haematoma was achieved in the following days, without further evidence of haemorrhagic complications. The patient was discharged on day 5 of admission and placed on dabigatran.

Figure 1 (A) Subcutaneous haematoma of the right leg with large haematic phlyctena; (B) detailed view of the lesion.

Despite decades of experience, warfarin still continues to be associated with haemorrhagic events.^{1–3} It is estimated that in 1 year, up to 6.5% of patients on anticoagulant therapy will experience a major bleeding event affecting their soft tissue, gastrointestinal tract or urinary tract.³ Approximately 1% of patients will develop a fatal bleed, often an intracranial haemorrhage.³ As far as the authors are concerned, this is the first warfarin-related spontaneous subcutaneous tissue haematoma of the lower limb reported in the literature. Clinicians should be aware of sporadic fluctuations of INR, even if the warfarin scheme is accomplished and regular INR control performed. Large haematomas such as that being reported can be life threatening, thus correction of the underlying coagulation abnormalities and close monitoring of the patient should be assured.^{1–3}

Learning points

- ▶ Despite prescription compliance and regular INR control, warfarin can be associated with spontaneous haemorrhagic events, even with moderately supratherapeutic INR.
- ▶ Correction of the coagulation disorder should be performed using fitomenadione, fresh frozen plasma or human prothrombin complex, with an individualised approach.
- ▶ Maintaining oral anticoagulation after a significant bleeding event is not consensual. However, novel oral anticoagulants could be a valid option after major bleeding, because of their shorter duration of action and new targeted antidotes.

Acknowledgements The authors would like to thank Dr Joana Alvarenga, Dr João Pereira and Dr Alberto Mello e Silva, for their contribution to this article.

Contributors All authors contributed equally to management of the patient as well as writing the article.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Roca B, Roca M. The new oral anticoagulants: reasonable alternatives to warfarin. *Cleve Clin J Med* 2015;82:847–54.
- 2 Tideman PA, Tirimacco R, St John A, et al. How to manage warfarin therapy. *Aust Prescr* 2015;38:44–8.
- 3 Zareh BS, Davis BS, Henderson MD. Reversal of warfarin-induced hemorrhage in the emergency department. *West J Emerg* 2011;12:386–92.

CrossMark

To cite: Pinheiro NC, Lopes A, Camões A, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-215134

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow