Pentazocine abuse in a healthcare professional

Aslı Enez Darçın,¹ Onur Cemal Noyan,² Serdar Nurmedov,³ Nesrin Dilbaz²

¹Istanbul Kanuni Sultan
Süleyman Eğitim ve Araştırma
Hastanesi, İstanbul, Turkey
²Uskudar University, İstanbul,
Uskudar, Turkey
³Acibadem Health Group,
İstanbul, Turkey

Correspondence to
Dr Onur Cemal Noyan,
onurnoyan@hotmail.com

Accepted 5 April 2016

DESCRIPTION

A 53-year-old man, a health professional from a Middle Eastern country, was admitted with parental overuse of pentazocine. He had first used pentazocine in the First Persian Gulf War for migraine. He began to abuse pentazocine, which he scaled up to 20 ampules per day and continued using for >15 years. Since 2006, because of political regulations in the healthcare system in his country, he obtained the drug from illegal sources instead of from pharmacies. He was abstinent for 9 months after a treatment programme. He suffered from musculoskeletal pain, diarrhoea and insomnia during early withdrawal.

He was using 10 30 mg pentazocine ampules intramuscularly in his left inguinal region for 6 months before admission. He had a wound in his left inguinal region that contained necrotic and infectious tissue (figure 1); he had been dressing this wound himself for months. He experienced guilty thoughts and was in a depressive mood.

He completed our 12-step therapy and was on quetiapine (200 mg), bupropion (150 mg) and naltrexone (50 mg) treatment after inpatient detoxification. He has been in follow-up in our outpatient clinic for 13 months and remains abstinent. He is functional in his occupation and social domains. After multiple surgeries in Turkey, his wound has healed.

Figure 1 Necrotic and infectious tissue in the inguinal region caused by chronic opioid injection.

Learning points

- ▶ Repeated injections of pentazocine may cause irregular ulcers with surrounding induration or hyperpigmentation. Pentazocine-induced ulcers can be an under-recognised cause of chronic non-healing ulcers.
- ▶ Rates of substance abuse among physicians are similar to those of the general population; physicians have no immunity to substance abuse disorders.
- ▶ The treatment outcomes for physicians with substance abuse disorders, when receiving appropriate assessment and treatment, can be considerably better than for the general population.

Repeated intramuscular injections of pentazocine, which is known to be abused, may cause soft tissue induration, fibrosis or ulceration, hyperpigmentation and myopathy.^{1 2} It is important to consider opioid injection in patients with a chronic wound, in the differential diagnosis. Although the rate of overall substance abuse is similar to that of the general population, abuse of opiates is more common in physicians. With appropriate treatment, many physicians with substance abuse disorders achieve long-term recovery and sobriety.³

Twitter Follow Onur Noyan at @iasonur

Contributors AED and SN conceived the idea for the manuscript and selected appropriate images. AED and OCN drafted the text. ND revised the text for important intellectual content. All the authors approved the final submitted version.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Mudrick C, Isaacs J, Frankenhoff J. Case report: injectable pentazocine abuse leading to necrotizing soft tissue infection and florid osteomyelitis. *Hand* 2011;6:457–9.
- 2 Goyal V, Chawla JM, Balhara YP, et al. Calcific myofibrosis due to pentazocine abuse: a case report. *J Med Case Rep* 2008;2:160.
- 3 Berge KH, Seppala MD, Schipper AM. Chemical dependency and the physician. *Mayo Clin Proc* 2009;84:625–31.

CrossMark

To cite: Darçın AE,
Noyan OC, Nurmedov S,
et al. *BMJ Case Rep*
Published online: [please
include Day Month Year]
doi:10.1136/bcr-2016-
215337

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow