Symmetrical digital gangrene after a high dose intravenous infusion of epinephrine and dopamine following resuscitation from cardiac arrest

Antonio Jose Reyes,¹ Kanterpersad Ramcharan,^{2,3} Patrick Harnarayan,⁴ Justin Mooteeram⁵

¹Neurology Unit, Department of Medicine, San Fernando Teaching Hospital, San Fernando, Trinidad and Tobago

²Department of Medicine, San Fernando Teaching Hospital, San Fernando, Trinidad and Tobago

³Department of Medicine, Surgi-Med Clinic, San Fernando, Trinidad and Tobago

⁴University of the West Indies, St Augustine, Trinidad and Tobago

⁵Department of Orthopedics, Sangre Grande Hospital, Sangre Grande, Trinidad and Tobago

Correspondence to

Dr Kanterpersad Ramcharan, kramcharan79@yahoo.com

Accepted 8 November 2016

DESCRIPTION

Symmetrical digital gangrene (SDG) is mostly due to low cardiac output and disseminated intravascular coagulation (DIC) (table 1).¹ We describe a 36-year-old right-handed man with SDG requiring amputation of 14 digits. He had received continuous intravenous infusion of epinephrine and dopamine given for a cardiac arrest that developed 1 hour after resection of a meningioma (table 2). In cardiogenic shock, he had decreased responsiveness, sinus tachycardia (135 bpm) and arterial hypotension (50/45 mm Hg). Advanced cardiac life support with mechanical ventilatory support was instituted. The administration of inotropic agents stabilised blood pressure to around 110/65, heart rate at 98 and mean arterial pressure (MAP) at 70 mm Hg. The patient's responsiveness improved gradually, but cardiovascular instability made him inotropic dependant.

On day 3 postcardiac arrest, acute gangrene of the tips of 10 toes and right 4 fingers developed. There was painful inflammatory oedema with erythematous reaction in the regions of demarcation between the normal and the gangrenous areas. All peripheral pulses were normal and symmetric on palpation. The digits were cold and pale with no capillary refill distally. No other relevant findings on physical examination were found. A colour Doppler ultrasound scan of the upper and lower limb vessels indicated normal blood flow. The patient recovery from brain surgery and cardiorespiratory instability were uneventful. Inotropic agents were gradually weaned off and eventually

discontinued. Mechanical ventilation support was discontinued successfully on day 7. Five tips of the toes underwent autoamputation over the next 3 months (figures 1A–E) after which the other affected distal phalanges of the digits were surgically removed, and the stumps were surgically repaired (figures 2A,B). The patient's digit stumps healed satisfactorily. Extensive medical investigations were conducted, and differentials were ruled out (table 1). Blood cultures, urine cultures and swabs cultures from the wounds were sterile.

Two years later, the patient had normal cardiovascular and musculoskeletal systems, normal nerve conduction studies of the limbs and duplex ultrasound of the upper and lower limb vessels indicated normal flow bilaterally. The patient has remained on social welfare due to multiple digital amputations.

Cases of digital necrosis and amputation have been associated with accidental epinephrine injections in fingers or interdigital regions (EpiPens) and intravenous inotropic agents.^{2–3} SDG has been reported at dopamine intravenous infusion >5.1 µg/kg/min in patients who had DIC and hypovolaemia. SDG can occur as an adverse event even though at therapeutic epinephrine dose infusion. Studies have shown the occlusion of small blood vessels when the intraluminal pressure falls below a critical value leads to compromised flow through digital arteries at perfusion pressures between 36 and 60 mm Hg.¹ We suggest this mechanism for the development of SDG in our patient. We believe severe hypotension may also have

Table 1 Causes of symmetrical digital gangrene

Hypercoagulable state	<i>Primary:</i> Antithrombin III deficiency, protein C and protein S deficiency, abnormalities of the fibrinolytic system and dysfibrinogenemias. <i>Secondary:</i> DIC, malignancy, pregnancy, use of oral contraceptives, myeloproliferative disorders, hyperlipidaemia, diabetes mellitus, sickle cell anaemia, polycythemia rubra vera and Waldenström's macroglobulinemia.
Ischaemic states	Low cardiac output, atherosclerosis, myocardial infarction.
Vasculitis	<i>Large vessels:</i> Polymyalgia rheumatica, Takayasu's arteritis and temporal arteritis. <i>Medium size vessels:</i> Buerger's disease, cutaneous vasculitis, Kawasaki disease, and polyarteritis nodosa. <i>Small vessels:</i> Behçet's disease, eosinophilic granulomatosis with polyangiitis, cutaneous vasculitis, Henoch Schönlein purpura, microscopic polyangiitis, granulomatosis with polyangiitis, Goller's vasculitis and cryoglobulinemia.
Autoimmune	Systemic lupus erythematosus, connective tissue disease, antiphospholipid syndrome and Raynaud's phenomenon.
Drugs	Dopamine, epinephrine and norepinephrine.
Infection-sepsis	Septic shock, leprosy.
Physical agents	Trauma, vibration, electrical injuries, burns, lightning strike, low temperature (frostbite) and high altitude.
Other causes	Congenital erythropoietic porphyria, ainhum, Lesch-Nyhan syndrome, syringomyelia and long-term tobacco smoking.
DIC, disseminated intravascular coagulation.

CrossMark

To cite: Reyes AJ, Ramcharan K, Harnarayan P, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-217977

Table 2 Drug therapy

Intravenous therapy	Dosage	Period of treatment
Epinephrine	1 mg intravenous push stat	Three times
Atropine	1 mg intravenous push stat	Three times
Epinephrine	10 mg/50 mL of solution 0.9% @ 8 mL/hour intravenous infusion	4 days then the drug was weaned down in a period of 3 days
Dopamine	400 mg/100 mL of solution 0.9% @ 10 mL/hour intravenous infusion	4 days then the drug was weaned down in a period of 3 days
Ceftriaxone	2 g intravenous daily	7 days
Subcutaneous drug	Dosage	Period of treatment
Enoxaparin	40 mg subcutaneous daily	10 days

Figure 1 (A) Wet gangrene of all five toes of the left foot. (B) Wet gangrene of all five toes of the right foot. (C) Right four fingers after surgical amputation with good stump healing. (D) Big toe autoamputation showing gangrenous phalanx. (E) Third toe autoamputation left foot.

contributed to the SDG in our patient. Appropriate dosage of intravenous inotropic agents to treat hypovolemic shock, cardiogenic shock or other life-threatening cardiovascular depression or instability should be individualised according to age, body mass index, MAP and cardiovascular status of the patient.

Figure 2 (A) Four right fingers stumps can be seen. (B) Ten toes stumps after 2 years are shown.

Learning points

- ▶ Intravenous epinephrine and/or dopamine and/or norepinephrine are used commonly. They can cause digital gangrene as an adverse drug reaction even at recommended dosage levels.
- ▶ Arterioles vasoconstriction seems to be the mechanism of gangrene induced by intravenous inotropic agents. Low cardiac output is also an important contributing factor to digital necrosis.
- ▶ Dosage of intravenous inotropic agents should be calculated individually and according to age, body mass index, mean arterial pressure (MAP) and cardiovascular condition, which may be safer than the usual standard of keeping MAP around 70 mm Hg.
- ▶ Digital gangrene has an increased risk of amputation of limbs with a decrease in functional capacity and quality of life.
- ▶ Since the differential diagnosis of digital gangrene is vast and no specific treatment for gangrene induced by intravenous inotropic agents has been determined, treatment has to be individualised.

Acknowledgements The authors thank the patient for allowing to use his medical data for teaching purposes. The authors also thank their other staff for caring for the patient. They are thankful to the photography department for preparing the images.

Contributors The conception, design and acquisition of data were carried out by AJR. The drafting of the article was also carried out by AJR. All authors contributed to the revisions for intellectual content and approve of the final version. KR and AJR are accountable for the article and will ensure that all questions regarding the accuracy and integrity of the article are investigated and resolved.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Colak T, Erdogan O, Yerebakan O, *et al.* Symmetrical peripheral gangrene and dopamine. *Ulus Travma Acil Cerrahi Derg* 2003;9:222–4.
- 2 Velissariou I, Cottrell S, Berry K, *et al.* Management of adrenaline (epinephrine) induced digital ischaemia in children after accidental injection from an EpiPen. *Emerg Med J* 2004;21:387–8.
- 3 Joynt G, Doedens L, Lipman J, *et al.* High-dose adrenaline with low systemic vascular resistance and symmetrical peripheral gangrene. *S Afr J Surg* 1996;34:99–101.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow