

# Frantz's tumour, solid pseudopapillary epithelial neoplasm (SPEN)

Orapin Tanapanpanit,<sup>1</sup> Samornmas Kanngurn,<sup>1</sup> Krit Pongpirul<sup>2,3</sup>


<sup>1</sup>Digestive Disease Center, Bumrungrad International Hospital, Bangkok, Thailand  
<sup>2</sup>Faculty of Medicine, Department of Preventive and Social Medicine, Chulalongkorn University, Bangkok, Thailand  
<sup>3</sup>Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

**Correspondence to**  
 Dr Krit Pongpirul, doctorkrit@gmail.com


Accepted 4 December 2016

## DESCRIPTION

A 29-year-old woman from Myanmar underwent medical check-up before having a baby. She was healthy and had no familial history of pancreatic disease or cancer. Physical examination revealed an ill-defined palpable mass arising from the left hypochondriac region. Blood tests including tumour marker levels were normal. Her abdominal ultrasound scan revealed a 1.37×1.28 inches mixed echogenic solid mass at distal pancreas without main pancreatic duct dilation (figure 1). Abdominal MRI demonstrated a well-encapsulated mass of size 1.70×1.61 inches, hyposignal T1-weighted (T1W) and heterogeneous hypersignal T2W masses with cystic and solid component with enhancement solid component at the pancreatic tail, with no evidence of intra-abdominal metastasis (figure 2). Distal pancreatectomy was performed. No adjuvant therapy was given, nor was splenectomy performed. She has been followed up as an outpatient for 2 years and has remained healthy without any sign of local recurrent or distal metastasis.



**Figure 1** Ultrasound scan finding: A solid mass like lesion at distal pancreas. No pancreatic duct dilation.




**Figure 2** MRI finding: A lobulated hyposignal T1-weighted (T1W), heterogeneous hypersignal T2W masses with cystic and solid component with enhancement solid component at pancreatic tail.


The mass had typical pathological features of solid pseudopapillary epithelial neoplasm (SPEN) (figures 3–5)—a combination of solid and cystic components with a cellular degenerative change



**Figure 3** Solid pseudopapillary neoplasm at distal pancreas.



**Figure 4** Pathological finding: Tumour tissue presenting cystic areas of haemorrhage and necrosis alternate with regions of solid growth, including notable encapsulation by pancreatic parenchyma (H&E, ×4).



**Figure 5** Pathological finding: Solid pseudo papillary structures composed of uniform cells surrounding small vessels (H&E, ×100).



**To cite:** Tanapanpanit O, Kanngurn S, Pongpirul K. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-218403

## Images in...

alternating with pseudopapillary formation along with uniform epithelioid cells with well-vascularised stroma and discohesive arrangement in the solid portions.<sup>1</sup> The differential diagnosis requires immunohistochemical staining. SPEN cells are usually absent when stains for endocrine markers are used and are characteristically positive for vimentin, CD10, CD56, and  $\alpha$ -1-antitrypsin.<sup>2</sup> The commonly encountered cystic neoplasms of the pancreas can be classified into five categories: serous microcystic adenoma, mucinous cystic neoplasms (cystadenoma or cystadenocarcinoma), intraductal papillary mucinous neoplasm (IPMN), cystic neuroendocrine neoplasms, and SPEN.

## Learning points

- ▶ Solid pseudopapillary epithelial neoplasm (SPEN) or Frantz's tumour is a rare benign condition, usually found in asymptomatic young woman who undergo screening imaging studies.
- ▶ Complete surgical resection is mandatory.

Among these uncommon pancreatic tumours, SPEN represents an exceedingly rare entity with low potential malignancy, usually found in asymptomatic young woman who undergo screening imaging studies.<sup>1 2</sup> It has many other names including Frantz tumour, as first described by Virginia Kneeland Frantz in 1959. Complete surgical resection is mandatory.<sup>2 3</sup>

**Contributors** OT was the primary physician, made the diagnosis, managed the case and prepared the first draft of the manuscript. SK provided pathological results and helped to draft the manuscript. KP contributed to the concept, and helped to draft and revised the manuscript.

**Competing interests** None declared.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- 1 Tang LH, Aydin H, Brennan MF, *et al*. Clinically aggressive solid pseudopapillary tumors of the pancreas: a report of two cases with components of undifferentiated carcinoma and a comparative clinicopathologic analysis of 34 conventional cases. *Am J Surg Pathol* 2005;29:512–19.
- 2 Madan AK, Weldon CB, Long WP, *et al*. Solid and papillary epithelial neoplasm of the pancreas. *J Surg Oncol* 2004;85:193–8.
- 3 Campanile M, Nicolas A, LeBel S, *et al*. Frantz's tumor: is mutilating surgery always justified in young patients? *Surg Oncol* 2011;20:121–5.

Copyright 2016 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact [consortiasales@bmjgroup.com](mailto:consortiasales@bmjgroup.com)

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow