OPEN ACCESS

CASE REPORT

Talus fracture in a 4-year-old child

Aurélien Michel-Traverso,¹ Trieu Hoai Nam Ngo,¹ Clémence Bruyere,² Marco Saglini³

¹Centre Hospitalier Universitaire
Vaudois, Lausanne, Switzerland
²Hopitaux Universitaires de
Geneve, Geneva, Switzerland
³EHF Morges, Morges,
Switzerland

Correspondence to
Dr Aurélien Michel-Traverso,
auremt@gmail.com

Accepted 24 March 2017

ABSTRACT

Fracture of the talus is uncommon in childhood. We report a case of talar neck fracture that occurred in a 4-year-old girl. We present the radiological findings, the orthopaedic follow-up and the clinical outcome.

BACKGROUND

Talus fracture is uncommon in childhood. A high level of clinical suspicion is needed to avoid missing this type of fracture, especially when it has a minimal displacement. Paediatric bone is more elastic than adult bone,¹ so fewer fractures occur in early life.

The prevalence for paediatric trauma in talus fracture is estimated to be five times rarer than for adult trauma.¹ The talar neck is the first site of fracture, followed by the talar body.^{2,3}

Potential bone remodelling for future growth is poor, which means avascular necrosis and osteoarthritis can occur if the fracture is not treated according to standard treatment methods.

CASE PRESENTATION

A 4-year-old girl presented to the emergency department with right foot pain after being hit by a heavy weight (metal wardrobe around 50 kg). Physical examination showed swelling, ecchymosis and tenderness on the dorsal side of the mid-foot. Foot and ankle mobilisation was painful, with

Figure 2 Day 0 lateral plain film.

inability to bear weight after receiving paracetamol and ibuprofen according to her weight. Additional examination was not possible due to the reduced compliance. X-rays (figures 1 and 2) confirmed the diagnosis of minimally displaced talar neck fracture⁴ (Hawkins type 1 neck of talus fracture without subtalar dislocation).

TREATMENT

The lesion was treated with a non-weight bearing short-leg cast for 4 weeks (figures 3 and 4) and in

Figure 1 Day 0 antero-posterior plain film.

Figure 3 1-Month antero-posterior plain film.

CrossMark

To cite: Michel-Traverso A,
Ngo THN, Bruyere C, et al.
BMJ Case Rep Published
Online First: [please
include Day Month Year].
doi:10.1136/bcr-2016-
215063

Figure 4 1-Month lateral plain film.

Figure 6 2-Month lateral plain film.

a weight-bearing synthetic cast for another 4 weeks. Analgesia with paracetamol and ibuprofen was given according to her weight; ice was applied on the cast for 10 min three times a day for the first 3 days to prevent swelling.

After 2 months the patient was pain free and had resumed all her activities. X-rays (figures 5 and 6) showed a consolidation of the fracture without evidence of avascular necrosis.⁵

OUTCOME AND FOLLOW-UP

Talar fracture is an uncommon pathology in children. For minimally displaced fractures, immobilisation with a cast has been

Learning points

- ▶ Talus fracture is an uncommon pathology in children which could be missed during diagnosis.
- ▶ Using a cast is a good treatment option with favourable outcomes for non-displaced and minimally displaced fractures.
- ▶ There is a need for long term follow-up in cases of talus fractures.

Figure 5 2-Month antero-posterior plain film.

proposed to avoid avascular necrosis and displacement of fracture.¹ In our case, the patient was treated conservatively. After 22 months, clinical findings revealed a favourable functional outcome. The mobilisation of the foot and ankle was complete without any pain.

DISCUSSION

Meier *et al*¹ studied the long-term outcome of neck or body talar fracture in 15 children (10 boys, 5 girls) with an average age of 10 years (4–16 years). All Hawkins types were studied with different treatment. Hawkins type 1 was treated by close reduction (if necessary) and with a cast immobilisation.

Patients with Hawkins type 1 and type 2 or body fractures achieved excellent results except two patients who had to be treated by an arthrodesis procedure. Smith *et al*⁶ studied 29 children with talus fractures sustained at an average age of 13.5 years. They observed no persistent osteonecrosis in patients younger than 12 years old and reported favourable outcomes in the majority of cases regardless of the mode of treatment.

The guidelines for the treatment of avascular necrosis and osteoarthritis caused by talus fracture in childhood trauma should be clearly defined, even if this rare type⁵ of fracture is not always recognised, especially with non-displaced fractures.

Contributors AM-T, T HNN and CB contributed to writing this case report. MS is the guarantor.

Competing interests None declared.

Patient consent Guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See:

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Meier R, Krettek C, Griensven M, *et al.* Fractures of the talus in the pediatric patient. *Foot and Ankle Surgery* 2005;11:5–10.
- 2 Byrne AM, Stephens M. Paediatric talus fracture. *BMJ Case Rep* 2012;2012:bcr1020115028.
- 3 Eberl R, Singer G, Schalamon J, *et al.* Fractures of the talus-differences between children and adolescents. *J Trauma* 2010;68:126–30.
- 4 Talkhani IS, Reidy D, Fogarty EE, *et al.* Avascular necrosis of the talus after a minimally displaced neck of talus fracture in a 6 year old child. *Injury* 2000;31:63–5.
- 5 Cartwright-Terry M, Pullen H. Non-operative management of a talar body fracture in a skeletally immature patient. *Acta Orthop Belg* 2008;74:137–40.
- 6 Smith JT, Curtis TA, Spencer S, *et al.* Complications of talus fractures in children. *J Pediatr Orthop* 2010;30:779–84.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow