Intraprocedural thrombus formation in a patient with anticoagulated atrial fibrillation

Sana Grover,¹ Krishan Kant Kataria,² Abraham G Kocheril³

¹Pt. B.D. Sharma Post Graduate Institute of Medical Sciences, Rohtak, Haryana, India

²Christie Clinic, Urbana, Illinois, USA

³Presence Covenant Medical Center, Urbana, Illinois, USA

Correspondence to

Dr Sana Grover,
sanagrover22@yahoo.com

Accepted 17 February 2017

DESCRIPTION

A 57-year-old man with 16-year history of persistent atrial fibrillation (AF) was admitted for electrophysiology study and catheter ablation. He was on amiodarone, carvedilol and warfarin. His international normalised ratio (INR) was 2.2. Intracardiac echocardiogram (ICE) prior to atrial septum puncture did not show any spontaneous echo contrast. After advancing the sheath across the interatrial septum, it was flushed with heparin. A few seconds after sheath placement and prior to placement of lasso catheter a linear thrombus developed which was visible on ICE (figure 1; videos 1 and 2). Activated clotting time (ACT) of >250 ms was maintained prior to trans-septal puncture and after thrombus was seen; heparin was administered to maintain ACT>300 ms. The sheath and catheter were retracted into the inferior vena cava, and with continuous heparin infusion, the thrombus resolved. During hospitalisation and

Video 2 A view of the left atrium showing the thrombus.

follow-up, patient remained asymptomatic without neurological complications.

Intraprocedural thrombus formation is a rare complication of AF ablation with incidence of 10%.¹ Strikingly, despite therapeutic INR, thrombus in our patient was extensive and formed within seconds after the trans-septal sheath was inserted into the left atrium (LA). Since the time across septum was <1 min, it is unlikely that the clot extruded from sheath. It is more likely that thrombus started forming outside sheath, as shown previously in another study.¹ This may be related to activation of coagulation cascade secondary to endothelial injury during catheter advancement into the LA.² Maintaining ACT>300 ms is routine part of our AF ablation procedures.

Around 90% of LA thrombi are formed in the LA appendage.³ Our case illustrates that thrombus can form elsewhere in the right atrium or LA.

Figure 1 Intracardiac echocardiogram image showing a thrombus strand extending through the interatrial septum.

Video 1 A view of left atrium showing a string of thrombus across the interatrial septum.

Learning points

- ▶ Atrial fibrillation can be highly thrombogenic.
- ▶ Although the left atrial appendage is the most common site of thrombus formation, the thrombus can form elsewhere in the right and left atrium.

Contributors AGK performed the procedure and conceived the idea of writing it up. SG and KKK did the literature search and wrote the manuscript. AGK did the proofreading and final editing.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

CrossMark

To cite: Grover S, Kataria KK, Kocheril AG. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-217527

REFERENCES

- 1 Ren J, Marchlinski F, Callans D. Left atrial thrombus associated with ablation for atrial fibrillation: identification with intracardiac echocardiography. *J Am Coll Cardiol* 2004;43:1861–7.
- 2 Watson T, Shantsila E, Lip GY. Mechanisms of thrombogenesis in atrial fibrillation: Virchow's triad revisited. *Lancet* 2009;373:155–66.
- 3 Odell JA, Blackshear JL, Davies E, et al. Thoracoscopic obliteration of the left atrial appendage: Potential for stroke reduction? *Ann Thorac Surg* 1996;61:565–9.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow