Alternative method of tractioning the Sengstaken-Blakemore tube

Evelyn Patricia Murphy,¹ Sean Michael O'Brien,² Mark Regan³

¹Department of Orthopaedics, University College Hospital Galway, Galway, Ireland
²Department of General Surgery, University College Hospital Galway, Galway, Ireland

³Department of GI/Surgery, Galway University Hospitals Group, Galway, Ireland

Correspondence to
 Dr Evelyn Patricia Murphy,
 evelynmurphy@hotmail.com

Accepted 4 March 2017

DESCRIPTION

Oesophageal varices commonly develop in the setting of portal hypertension. A Sengstaken-Blakemore (SB) tube is sometimes emergently required to control oesophageal bleeding.^{1,2}

The normal portal pressure varies from 5 to 10 mm Hg. Portal hypertension occurs when the portal pressure rises above 10 mm Hg. Pressures can rise up to 30 mm Hg, so the pressure exerted and maintained on the oesophageal balloon to tamponade the varices will need to be above 30 mm Hg.

The current practice is to attach a 1 L bag of saline to the tube to maintain traction. However, there are risks associated with this method of traction. These risks include pharyngeal ulcers and mouth ulcers from the local pressure effect. We describe an alternative method of maintaining traction on the SB tube.

We describe the practice of securing the tube with a pair of wooden tongue depressors and padding as shown in figure 1. This has been used by the lead author for 25 years with no adverse events. The patient population includes patients with oesophageal varices. The patients ranged from 18 to 85 years of age. Over 50 patients have been treated by the lead author using this method.

This is an alternative safe method of applying traction without the adverse local pressure effects.

Equipment needed

Equipment needed are McGill forceps, lubrication jelly, 2×60 cc syringes with a luer lock, one large bladder irrigation syringe, a sphygmomanometer or any other hand-held device for measuring pressure, and contrast with water for insertion into the gastric balloon.

Insertion

1. These patients are usually intubated as the risk of airway contamination is extremely high if balloon tamponade is required to control bleeding. If possible, these cases should be performed in theatre.
2. Check and confirm there are no air leaks in the gastric and oesophageal balloons.
3. The tube has markings on it; make sure that these markings are visible when inserting the tube. When it is being inserted endoscopically, direct visualisation of the gastro-oesophageal junction is possible.
4. Insert a mix of contrast dye and water into the gastric balloon. This will enable visualisation on the subsequent chest X-ray to confirm position postprocedure. Once 150–200 mL of water/contrast mix should have been filled; close the aspiration port.

Figure 1 Traction device in use on a patient.

To cite: Murphy EP, O'Brien SM, Regan M. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2016-218401

5. Pull the balloon until it rests firmly against the gastro-oesophageal junction. An assistant is necessary to maintain this traction until the definitive traction has been set up.
6. Insert air into the oesophageal balloon. The author prefers to aim for a pressure of 28–30 mm Hg. The normal portal pressure is 5–10 mm Hg and the consequences of pressure necrosis in the oesophagus are serious. The oesophageal balloon is routinely inflated in all patients for a minimum of 4 hours and then the situation is re-evaluated. The oesophageal balloon is deflated at the 4 hour mark and then can be reinflated if necessary. The traction device has been attached with the oesophageal and gastric balloon inflated.

Maintain traction

Various methods of maintaining traction have been described in the literature to date. Some of the more common methods include attaching a 500–1000 mL bag of saline to the SB tube. Other options occasionally favoured in the USA include securing the SB tube to the rim of a helmet.³ However, several users have described this as cumbersome in terms of nursing care. Other institutions have described a method of splitting a tennis ball in half to maintain traction and fix it to the patients face. However, this has the disadvantage of a pressure effect on the patient's teeth and lips. Tennis balls are not readily available to most institutions in an emergent situation. Record the marking at the teeth. Documentation will ensure the tube does not slip and that traction is maintained.

Set up of alternative device

Step 1: Secure a thin rolled up 10×10 cm gauze to the longitudinal surface of each tongue depressor. A sleek tape may be used to secure the gauze (figure 2).

Step 2: Wrap the tape around the cranial end of the SB tube using two twists, do not cut it from the roll. The SB tube should be cleaned with a swab prior to ensure it is dry before (figure 3).

Step 3: Wrap the sleek tape around both sides of the tongue depressor construct and then finish by wrapping a layer circumferentially around the SB tube (figure 3).

Figure 2 Demonstration of adhesive tape around the tongue depressors.

Figure 3 Demonstration of adhesive tape attachment to the Sengstaken-Blakemore tube.

Step 4: Apply a foam wrap to the surface which is in contact with the mouth and lips. We use the same foam that is used in casts as this has been designed to withstand considerable friction (figure 4).

Step 5: Take note of the marking and educate staff on the construct. This may be repositioned by the nursing staff by twisting/altering the direction.

Aftercare

Regular suctioning of the oesophageal and gastric lumen is necessary.

Frequent oropharyngeal suctioning is necessary.

Antibiotic prophylaxis and 48 hours of terlipressin are required.

Repeat endoscopy at 24 hours is advised.

Figure 4 Demonstration of the Sengstaken-Blakemore tube with foam.

Learning points

- ▶ The combination of this traction method and ensuring a relatively low pressure is maintained in the oesophageal balloon has resulted in few/no pressure-related complications.
- ▶ There has been a positive experience with this method. It is cost-effective, easy to assemble and does not have the same rate of pressure-related ulcers as reported in the literature with the saline bag.
- ▶ There has been a lack of trials or studies reporting on the differing traction methods in securing the Sengstaken-Blakemore (SB) tube. Limitations could include a lack of a standardised method to secure the SB tube.

In our unit, the oesophageal balloon is deflated at 6–12 hours postinflation.

Repeat endoscopy should be carried out if possible in theatre. These patients are usually intubated.

Potential limitations of using this method could include unfamiliarity with the device. However, this would be overcome by education.

Contributors EPM used the device. SMOB proof read the manuscript and performed literature review. MR devised the device and proof read the manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Helmy A, Hayes PC. Review article: current endoscopic therapeutic options in the management of variceal bleeding. *Aliment Pharmacol Ther* 2001;15:575–94.
- 2 Seet E, Beevee S, Cheng A, *et al*. The Sengstaken-Blakemore tube: uses and abuses. *Singapore Med J* 2008;49:e195–7.
- 3 Kashiwagi H, Shikano S, Yamamoto O, *et al*. Technique for positioning the Sengstaken-Blakemore tube as comfortably as possible. *Surg Gynecol Obstet* 1991;172:63.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow