Transabdominal wall lipoma

Amari T Thompson, Samuel J Ford, Anant Desai, David Gourevitch

Queen Elizabeth Hospital
Birmingham, Birmingham, UK

Correspondence to
Dr Amari T Thompson,
amari.thompson@doctors.
org.uk

Accepted 17 March 2017

DESCRIPTION

A 60-year-old man presented with a painful, enlarged palpable mass in the left iliac fossa. A CT scan of the abdomen and pelvis demonstrated a large heterogeneous mass extending through the abdominal wall musculature (figure 1), which was suspected to be an atypical lipomatous tumour (ALT) or well-differentiated liposarcoma (WD-LPS).

Intraoperatively, a large firm lipomatous mass was noted in close proximity to the left spermatic cord. It extended into the retroperitoneal space and transgressed the internal oblique to lie deep to the external oblique aponeurosis, adopting an unusual 'dumbbell' shape. Due to clinical uncertainty as to whether the mass was simply in close proximity to the spermatic cord or a cord liposarcoma, a left radical orchidec-tomy was performed en bloc with the tumour.

Lipomas are the most common soft tissue mesenchymal tumour,¹ and are histologically benign. It is highly unusual for them to pass through tissue planes. Lipomas lack MDM2 amplification on cyto-genetic testing; unlike ALTs and WD-LPSs which are locally progressive, have a propensity to recur and require specialist management. The terms ALT and WD-LPS are synonymous as these tumours are

Figure 1 Lipoma passing through the abdominal wall musculature.

Learning points

- ▶ Lipomas are the most common soft tissue mesenchymal tumour and are typically <10 cm in diameter.
- ▶ The terms well-differentiated liposarcoma and atypical lipoma are synonymous.
- ▶ Lipomatous lesions which are >10 cm in the extremity, intramuscular or deep seated must be referred to a specialist unit for management.

morphologically and karyotypically identical.¹ ALT describes lesions in superficial sites and extremities² which are more amenable to complete surgical resection with a wide margin. WD-LPS describes lesions arising in the retroperitoneum, mediastinum and deep pelvis—locations that are less amenable to radical excision due to surrounding structures. Nagano *et al*² have devised a scoring system to differentiate lipomas and ALTs using MRI characteristics: diameter, depth, septa and fat saturation enhancement. The system appears promising, but requires further validation.

Contributors The patient was admitted under SJF who was assisted by ATT in drafting the article. AD and DG revised the article and were also heavily involved in managing the patient.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Fletcher CDM, Unni KK, Mertens F. *World Health Organization classification of tumours. Pathology and genetics of tumours of soft tissue and bone*. Lyon: IARC Press, 2002.
- 2 Nagano S, Yokouchi M, Setoguchi T, *et al*. Differentiation of lipoma and atypical lipomatous tumor by a scoring system: implication of increased vascularity on pathogenesis of liposarcoma. *BMC Musculoskelet Disord* 2015;16:36.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow

CrossMark

To cite: Thompson AT, Ford SJ, Desai A, *et al*. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2016-218775