Trousseau syndrome

Tatsuya Fujikawa,¹ Yasuharu Tokuda²

¹Mitoyo General Hospital, Kan-onji, Japan

²Department of General Internal Medicine, Tsukuba University, Mito, Japan

Correspondence to
Dr Tatsuya Fujikawa,
tfujikawa-gi@umin.ac.jp

Accepted 23 February 2017

DESCRIPTION

A 45-year-old woman was admitted to our hospital because of impaired consciousness. She had a huge tumour measuring 10 inches in diameter on the left breast (figure 1A). Although the tumour had been increasing in size since 3 years, the patient had adamantly refused to visit a hospital. She had no history of hypertension or atrial fibrillation. Symmetrical high-intensity lesions in the thalamus were observed on MRI (diffusion-weighted images) (figure 1B, arrow). Enhanced chest CT imaging revealed a large tumour on the left breast (figure 1C, arrowheads) with metastases in the lymph nodes and lungs (figure 1C, arrows) as well as cardiac invasion. Breast mass biopsy revealed the tumour to be a mucinous carcinoma. Impaired consciousness was considered to be caused by bilateral paramedian thalamic infarction.

Our case did not have common risk factors for atherosclerosis such as hypertension, diabetes and smoking. The patient was diagnosed with Trousseau syndrome, a state of hypercoagulability associated with the mucinous breast carcinoma. She regained consciousness after 3 days of conservative therapy for cerebral infarction but refused chemotherapy and received end-of-life care at home.

Trousseau syndrome comprises thromboembolic disorders in arterial or venous systems associated with malignancy, especially mucin-secreting carcinomas of the pancreas and gastrointestinal tract, lung cancer and ovarian cancer.^{1 2} Thromboembolism in the lower extremities, lung, brain, heart, kidneys or spleen may lead to death among patients with this syndrome.^{1 3} The syndrome may be treated by curative tumour resection and anticoagulation therapy,^{1 3} but when patients present with advanced cancer, as that in our case, curative therapy cannot be used. Trousseau syndrome with cerebral infarction should be considered to be a cause of impaired consciousness in patients with advanced cancer, particularly mucinous carcinoma.

Learning points

- ▶ Trousseau syndrome is a hypercoagulability syndrome associated with cancer.
- ▶ Mucinous carcinoma is one of the most common cancers that is likely to cause hypercoagulability.
- ▶ Trousseau syndrome with cerebral infarction should be considered to be a cause of impaired consciousness in patients with advanced cancer.

Figure 1 (A) A huge tumour measuring 10×10 inches in diameter with pigimentary changes was observed on the left breast. (B) MRI revealed symmetrical high-intensity lesions in the thalamus on diffusion-weighted images (arrow). (C) Enhanced CT revealed a large tumour on the left breast (arrowheads) with lung metastases (arrows).

CrossMark

To cite: Fujikawa T, Tokuda Y. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2017-219249

Contributors TF wrote manuscript and contributed to discussion. YT reviewed and edited the manuscript and contributed to discussion.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Chen YL, Lin KH, Lin MC, *et al.* Malignant struma ovarii complicated by Trousseau's syndrome and repeated episodes of cerebral ischemic strokes: a case report. *Gynecol Oncol Case Rep* 2011;2:35–8.
- 2 Caine GJ, Stonelake PS, Lip GY, *et al.* The hypercoagulable state of malignancy: pathogenesis and current debate. *Neoplasia* 2002;4:465–73.
- 3 Nakayama M, Iha T, Kanazawa K. Unusual Trousseau's syndrome in ovarian carcinosarcoma: multiple systemic thromboembolic events. *J Obstet Gynaecol* 2002;22:699–700.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow