Giant hiatal hernia: beware of the supine ICU chest X-ray!

Vincent Bunel,^{1,2} Pierre Mordant,^{2,3} Lara Ribeiro,⁴ Bruno Crestani^{1,2}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bcr-2017-219668>)

¹Bichat Hospital, Department of Pneumology A, DHU Fire, Assistance Publique - Hopitaux de Paris, Paris, France

²Universite Paris Diderot UFR de Medecine Site Xavier-Bichat, Paris, France

³Bichat Hospital, Department of Vascular and Thoracic Surgery, Assistance Publique - Hopitaux de Paris, Paris, France

⁴Bichat Hospital, Department of General and Digestive Surgery, Assistance Publique - Hopitaux de Paris, Paris, France

Correspondence to
Dr Vincent Bunel,
vincent.bunel@gmail.com

Accepted 15 March 2017

DESCRIPTION

An 85-year old woman with a medical history of mild hiatal hernia was admitted to the medical intensive care unit for respiratory distress associated with hypoxaemia and hypercapnia. A diagnosis of cardiogenic oedema was suspected. The situation slightly improved after high-dose diuretics and non-invasive ventilation. The anteroposterior supine chest X-ray revealed a right thoracic opacity. The patient was transferred to the pulmonology unit. A classical erected posteroanterior chest X-ray revealed a large air-fluid level in the right hemithorax (figure 1). An hydropneumothorax or a lung abscess was suspected. A chest CT scan revealed a giant hiatal hernia containing the stomach, the first duodenum, the spleen, the caudal part of the pancreas and the left colic flexure, all elevated in the right hemithorax (figure 2A–C; video 1). A laparoscopic surgical procedure was performed which involved the excision of the hernia sac, a suture repair of crural pillars, with oesophagogastropexy and an antireflux valve creation (figure 2D–F).

Figure 1 Posteroanterior chest X-ray showing large right chest opacity with air-fluid level on the right part of the thorax. The right hemidiaphragm is not visible.

Figure 2 CT scan before (A,B,C) and after (D,E,F) surgery. The stomach (A,E, triangle), left colonic flexure (A,F, arrow) and spleen (C,E, star) are visible in the hernia sac and in their right position after surgery; the liver is marked by a cross (B,C,D,E). Postoperative CT images show moderate postoperative pleural effusion (D) and significant diminution of hiatal hernia (F).

The postoperative period was uneventful and the patient was discharged on postoperative day 13. Comparison of the preoperative and postoperative work-up showed a great improvement in dyspnea (Modified Medical Research Council (mMRC) Dyspnea Scale 2 vs mMRC 4 before surgery), room air oxygen saturation ($\text{SpO}_2=97\%$ vs 90%) and lung function (forced vital capacity= 107% predicted vs 48% predicted). Huge hiatal hernia is particularly observed in elderly, and may cause dyspnoea¹ and cardiac failure due to heart compression.² Surgical repair is usually associated with a significant improvement of both symptoms and in pulmonary function tests.³

Contributors Conception and design: VB, BC. Analysis and interpretation: VB, PM, LR, BC. Drafting the manuscript for important intellectual content: VB, PM, BC.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed

To cite: Bunel V, Mordant P, Ribeiro L, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-219668

Learning points

- Beware of anteroposterior supine chest X-ray.
- Think of hiatal hernia for all hydro-aeric opacity in the thorax, even on the right side.
- Giant hiatal hernia surgery can improve dyspnoea.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Sahin C, Akin F, Cullu N, *et al.* A large Intra-Abdominal hiatal hernia as a rare cause of dyspnea. *Case Rep Cardiol* 2015;2015:1–4.
- 2 Matar A, Mroue J, Camporesi E, *et al.* Large hiatal hernia compressing the heart. *Am J Cardiol* 2016;117:483–4.
- 3 Carrott PW, Hong J, Kuppusamy M, *et al.* Repair of giant paraesophageal hernias routinely produces improvement in respiratory function. *J Thorac Cardiovasc Surg* 2012;143:398–404.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow