Truth will out: a rare case of amyloid cardiomyopathy

Jesse Jongman, Suresh Ghoerbien, Marco Götte

Department of Cardiology,
HagaZiekenhuis, Den Haag, The
Netherlands

Correspondence to
Dr Jesse Jongman,
jessejongman@hotmail.com

Accepted 28 March 2017

DESCRIPTION

A 65-year-old patient with no medical history of notice was referred to our cardiology outpatient clinic for the evaluation of exertional dyspnoea. After a thorough work-up, the suspicion of an amyloid cardiomyopathy was raised by the cardiac magnetic resonance (CMR) findings (figure 1A–C), but laboratory tests, abdominal fat pad and bone marrow biopsies failed to prove the diagnosis. Eventually endomyocardial biopsies confirmed the diagnosis of secondary (AA) amyloidosis. Furthermore, the CMR scout images unveiled a large abdominal mass (figure 1D), with subsequent CT being suggestive of a pancreatic neuroendocrine tumour with multiple hepatic metastases.

Clinically apparent heart disease or cardiac amyloidosis is uncommon in secondary amyloidosis.¹ Secondary amyloidosis may complicate chronic inflammatory diseases and is rarely associated with other causes, for example, neoplasms.¹ In the diagnosis of cardiac amyloidosis, the characteristic echocardiographic finding of an increased echogenicity of the myocardium has limited

specificity and sensitivity.² In contrast, CMR imaging with a distinctive late gadolinium enhancement pattern can provide evidence strongly suggestive of cardiac amyloid deposition and can be used to guide myocardial biopsies, which have a higher sensitivity compared with subcutaneous fat or rectal biopsies.^{2,3}

In our patient, the characteristic CMR findings proved to be pivotal for the diagnosis of the secondary or AA amyloidosis with cardiac manifestation. Furthermore, the CMR scout images unveiled an important incidental finding, that is, a large abdominal mass, possibly associated with the amyloidosis in this patient.¹

Learning points

- ▶ Cardiac magnetic resonance (CMR) has a vital role in the work-up of suspected (cardiac) storage diseases.
- ▶ Endomyocardial biopsy has a high sensitivity for the diagnosis of cardiac amyloidosis.
- ▶ Use all CMR images, including the scout images.

Figure 1 Cardiac magnetic resonance images showing diffuse subendocardial late gadolinium enhancement of the left ventricle, right ventricle and atria. (A) Four-chamber image; (B) two-chamber image; (C) midventricular, short-axis image and (D) balanced steady-state free precession (SSFP) scout image showing a large abdominal mass.

Contributors JJ is the first author and wrote the manuscript. SG helped with the interpretation of the CMR and supplied the images. MG supervised the CMR reporting and supervised the clinical course of the patient as well as the manuscript.

Competing interests None declared.

Patient consent Obtained from next of kin.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Lachmann HJ, Goodman HJ, Gilbertson JA, et al. Natural history and outcome in systemic AA amyloidosis. *N Engl J Med* 2007;356:2361.
2. Austin BA, Tang WH, Rodriguez ER, et al. Delayed hyper-enhancement magnetic resonance imaging provides incremental diagnostic and prognostic utility in suspected cardiac amyloidosis. *JACC Cardiovasc Imaging* 2009;2:1369.
3. Pellikka PA, Holmes DR, Edwards WD, et al. Endomyocardial biopsy in 30 patients with primary amyloidosis and suspected cardiac involvement. *Arch Intern Med* 1988;148:662.

CrossMark

To cite: Jongman J, Ghoerbien S, Götte M. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-219775

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow