Occult femoral neck fracture

Naoki Yoshida,¹ Yohei Komaru,² Yoshihiko Tsuchida¹

¹Department of Orthopedic Trauma Surgery, Shonan Kamakura General Hospital, Kanagawa, Japan

²Nephrology, Immunology and Vascular Medicine, Shonan Kamakura General Hospital, Kanagawa, Japan

Correspondence to

Dr Naoki Yoshida,
nyoshida.thk@gmail.com

Accepted 16 March 2017

DESCRIPTION

A 77-year-old woman presented to the emergency department with persistent pain that developed in the right hip after a fall from a standing height. A radiograph and CT of the right hip revealed no evidence of fracture (figure 1A,B), but MRI revealed oedema of the right femoral neck (T1-weighted

Figure 1 (A) A radiograph of the right hip revealed no evidence of fracture. (B) A CT of the right hip revealed no evidence of fracture. (C) MRI (T1-weighted coronal imaging) revealed oedema of the right femoral neck. (D) Patient was treated with osteosynthesis.

Learning points

- ▶ Occult femoral neck fractures may not be easily diagnosed on a radiograph.
- ▶ If occult femoral neck fracture is misdiagnosed, the non-displaced fracture may become displaced. In this case, a patient may require treatment with hip arthroplasty.
- ▶ MRI is one of the best tools for diagnosing occult fractures.

coronal imaging; figure 1C). She was treated with osteosynthesis (figure 1D) to stabilise the femoral neck fracture, which resulted in a good functional outcome at 6-month follow-up.

Occult femoral neck fractures are frequently misdiagnosed.¹ Once misdiagnosed, the non-displaced fracture may become displaced. When this occurs, patients may need to be treated with hip arthroplasty, which is more invasive than osteosynthesis.

In the context of persistent hip pain where plain radiographs and CT reveal no abnormality, consideration should be given to the use of MRI as it can be more accurate at detecting occult hip fractures.²

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement The patient's care was overseen by NY, YT, YK, YT provided assistance with the drafting of the manuscript. All authors approved the manuscript prior to submission.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Krestan CR, Nemec U, Nemec S. Imaging of insufficiency fractures. *Semin Musculoskelet Radiol* 2011;15:198–207.
- 2 Lim KB, Eng AK, Chng SM, et al. Limited magnetic resonance imaging (MRI) and the occult hip fracture. *Ann Acad Med Singapore* 2002;31:607–10.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow

CrossMark

To cite: Yoshida N, Komaru Y, Tsuchida Y. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-219868