Scalp vein sign: a forgotten clinical clue?

Shriraam Mahadevan,^{1,2} Adlyne Reena Asirvatham,³ Subramanian Kannan,⁴ Dhalapathy Sadacharan^{5,6}

¹Endocrinology Department, Sri Ramachandra Medical College, Chennai, India

²Endocrinology, Endocrine & Speciality Clinic, Chennai, India

³Sri Ramachandra University Medical College, Chennai, India

⁴Endocrinology, Diabetes and Bariatric Medicine, Narayana Health City, Bengaluru, India

⁵Endocrine Surgery, Madras Medical College, Chennai, India

⁶Department of Endocrine Surgery, Endocrine and Speciality Clinic, Chennai, India

Correspondence to

Dr Shirraam Mahadevan, mshriraam@gmail.com

Accepted 3 April 2017

DESCRIPTION

A 72-year-old man presented for general health check. He was asymptomatic except for hard of hearing and uses hearing aid for the last 4 years. Clinical examination revealed very prominent tortuous superficial temporal veins bilaterally, more on the left side (figure 1) with no bruit and a warm non-tender skull on palpation. Serum chemistries showed significantly elevated alkaline phosphatase 1520 U/L (normal, 45–140 U/L) with normal liver, kidney and haematological parameters. Serum calcium, phosphorus, 25-hydroxy vitamin D and parathyroid hormone levels were normal. Skull radiograph showed moth-eaten appearance suggestive of Pagetoid bone (figure 2). With this clinico-biochemical picture in an otherwise asymptomatic individual, Paget's disease of bone was strongly considered. It was confirmed by significant uptake of technetium-methylene diphosphonate (MDP) in skull and right hemipelvis. He was managed with intravenous zoledronic acid, calcium and vitamin D supplementation. Serum alkaline phosphatase normalised at 6 months follow-up, though hearing and scalp vein distension did not improve significantly. Paget's disease of bone, hitherto considered to be rare, is now being increasingly diagnosed in India.¹ Skull involvement is commonly seen in polyostotic

Figure 2 Lateral view of skull radiograph showing the typical 'moth-eaten' appearance of Pagetoid bone.

Paget's disease. Prominent superficial temporal vein which occurs due to increased vascularity and arteriovenous shunting was earlier described as 'Scalp vein sign'.² In a case series of 290 patients, this sign was observed in 45% of those with skull involvement³ and hence may serve as a useful clinical clue in asymptomatic individuals. Conditions which could mimic this sign include giant cell arteritis and metastases of follicular thyroid carcinoma, wherein the arteries become more prominent and tender.

Learning points

- ▶ Paget's disease of bone is not uncommon in South India.
- ▶ Skull involvement is very common in polyostotic Paget's disease.
- ▶ Prominent superficial temporal vein is a simple clinical sign that along with raised alkaline phosphatase may aid in diagnosis of asymptomatic Paget's disease of bone.

Contributors SM was involved in the diagnosis and management of the patient, literature search and preparation of the manuscript. ARA was involved in the diagnosis, literature search and manuscript review. SK was involved in patient diagnosis and critical review of the manuscript. DS was involved in manuscript review and critical comments.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Anjali S, Thomas N, Rajaratnam S, et al. Paget's disease of bone: experience from a centre in southern India. *J Assoc Physicians India* 2006;54:525–9.

Figure 1 Clinical picture showing prominent, tortuous, left superficial temporal vein with hearing aid.

CrossMark

To cite: Mahadevan S, Asirvatham AR, Kannan S, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-220259

2 Altman RD, Gray RG. Scalp vein sign in Paget's disease of bone. *J Rheumatol* 1982;9:624–6.

3 Altman RD. Musculoskeletal manifestations of Paget's disease of bone. *Arthritis Rheum* 1980;23:1121–7.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow