Anal herpes

Andre Goulart,^{1,2} Jose Pinto,¹ Pedro Leão^{1,2}

¹General Surgery, Hospital de Braga, Braga, Portugal

²School of Health Sciences, University of Minho, Braga, Portugal

Correspondence to

Dr Andre Goulart, andre.b.goulart@gmail.com

Accepted 7 June 2017

DESCRIPTION

A 27-year-old man without significant past medical history presented to the emergency department with a 5-day evolution of abdominal and anal pain and haematochezia. Physical examination revealed perianal erythema and a painful digital anal examination without other findings. CT scans showed signs of proctitis (slight thickening of rectal wall and slight stranding of perirectal fat). Flexible rectosigmoidoscopy didn't reveal any pathological abnormalities. The patient was discharged with antibiotics and analgesics with the suspicion of proctitis. Three days later, the patient returned to the emergency department with worsening of the anal pain that irradiated to the buttocks and the physical examination revealed red bumps with ulceration (figure 1). Anal herpes was suspected and the patient was treated with an antiviral (brivudine) and carbamazepine for symptomatic relief. Seven days after treatment the patient had complete resolution of his symptoms and there were no cutaneous lesions.

Herpes simplex virus (HSV) infections are the main cause of genital ulcers worldwide, and HSV-2 is the serotype most frequently implicated in its aetiology.¹ A visible outbreak consists of single or clustered vesicles on the genitalia, perineum, buttocks, upper thighs or perianal areas that ulcerate before resolving.² Antiviral agents are effective in controlling clinical episodes, but do not eradicate infection, which remains latent for the life of a patient.³

Learning points

- ▶ Extragenital herpes is a difficult diagnosis.
- ▶ High levels of suspicion and careful history-taking and physical examination are required to make a correct diagnosis.

Contributors AG, JP and PL contributed equally in planning, writing and reviewing the article.

Competing interests None declared.

Patient consent Obtained.

Figure 1 Anal herpes.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

1. Gupta R, Warren T, Wald A, *et al.* Genital herpes. *Lancet* 2007;370:2127–37.
2. Groves MJ. Genital herpes: a review. *Am Fam Physician* 2016;93:928–34.
3. Garland SM, Steben M. Genital herpes. *Best Pract Res Clin Obstet Gynaecol* 2014;28:1098–110.

CrossMark

To cite: Goulart A, Pinto J, Leão P. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-220856

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow