Extensive cutaneous involvement due to herpes simplex virus infection

Ana Vilaça,¹ Teresa Pimentel²

¹Internal Medicine, Hospital de Braga, Braga, Portugal
²Internal Medicine, Hospital de Braga, Braga, Portugal

Correspondence to

Dr Ana Vilaça,
 ana.vilaca.rodrigues@gmail.com

Accepted 23 May 2017

DESCRIPTION

A 39-year-old woman, with a medical history of oligofrenia, obesity and varicella at 8 years of age, presented to the emergency department (ED) with multiple skin lesions of upper limb, which began with the appearance of vesicles, associated with intense pain and pruritus and with 3 days of evolution. No fever was reported. She was discharged home medicated with acyclovir and hydroxyzine.

Three days later, she returned to the ED with worsening complaints of pain and itching, and extension of cutaneous lesions throughout the body.

On physical examination, she was febrile (T: 38.5°C), with erythematous-pruriginous lesions, some of which were typically targeted, associated with numerous bullae dispersed throughout the body with oral mucosa involvement (figure 1A–C).

Laboratory investigation revealed normal complete blood count. Serology tests confirmed that the patient was positive for herpes simplex virus (HSV) and there was threefold rise in antibody titre. Serology for other infectious agents, microbiology and autoimmune blood work studies were negative.

At this point, extensive cutaneous involvement due to HSV infection was assumed as a diagnosis.

Learning points

- ▶ HSV infections are the most frequent within the herpes virus group and generally have a benign and self-limiting course. However, its presentation may be atypical in some cases, manifesting itself as a widespread infection and, rarely, life-threatening.^{1,2}
- ▶ Reactivation of HSV after childhood may occur spontaneously or through a state of immunosuppression. In fact, the more exuberant manifestations occur more frequently in immunocompromised patients.² However, the authors emphasise the exuberance of cutaneous lesions that can occur in an immunocompetent patient, as we reported in our case.
- ▶ Given these cutaneous lesions, we have to keep in mind differential diagnoses that manifest in a similar way as, for example, autoimmune blistering disorders.³ Thus, the authors highlight the importance of clinical awareness of these atypical manifestations, so that we can arrive at a correct diagnosis with the least delay.

Figure 1 (A) Erythematous lesions associated with numerous bullae dispersed throughout the right upper limb. (B) Erythematous lesions associated with numerous bullae dispersed throughout the left upper limb. (C) Erythematous lesions some of which were typically targeted in abdomen.

Figure 2 (A) Erythematous lesions in healing phase (right upper limb). (B) Erythematous lesions in healing phase (left upper limb). (C) Erythematous and target lesions in healing phase (abdomen).

CrossMark

To cite: Vilaça A, Pimentel T.
BMJ Case Rep Published
 Online First: [please
 include Day Month Year].
 doi:10.1136/bcr-2017-
 221003

The patient was treated with a 7-day course of acyclovir (1000 mg/day) and with topical corticosteroid. She also was medicated with hydroxyzine (25 mg/day), while she had itching complaints.

After 1 week, the patient presented significant improvement of the lesions (figure 2A–C).

Contributors AV: Substantial contributions to conception and design, acquisition of data, analysis and interpretation of data, drafting the article, critical revision of the article and final approval of the version to be published. TP: Conception and design, analysis and interpretation of data, critical revision of the article and final approval of the version to be published.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Drago F, Rebora A. Viral reactivation and skin eruptions. *Dermatology* 2003;207:1–2.
- 2 Drago F, Ciccarese G, Gasparini G, *et al.* Contemporary infectious exanthems: an update. *Future Microbiol* 2017;12:171–93.
- 3 Lecluse AL, Bruijnzeel-Koomen CA. Herpes simplex virus infection mimicking bullous disease in an immunocompromised patient. *Case Rep Dermatol* 2010;2:99–102.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow