The telescoping aorta: a case of circumferential aortic dissection with intimo-intimal intussusception into the left ventricle

Majd Ibrahim, John Rashid, Stephen Craig Kurtz, Sudhir Mungee

Department of Internal Medicine/Division of Cardiology, University of Illinois College of Medicine at Peoria/OSF Saint Francis Medical Center, Peoria, Illinois, USA

Correspondence to

Dr Majd Ibrahim,
ibrahim_majd@yahoo.com

Accepted 16 August 2017

DESCRIPTION

A 67-year-old man with history of tobacco smoking presents to the emergency department with sudden onset, retrosternal chest pain radiating to the neck and associated with diaphoresis and nausea. On examination, his blood pressure was 106/42 mm Hg, pulse of 73 beat/min. He was pale, diaphoretic with IV/VI diastolic blowing murmur of unknown chronicity heard at the left sternal border. His distal pulses were faint but symmetric in all extremities. No marfanoid features. ECG showed sinus rhythm with ST segment elevation in leads

Figure 1 Admission ECG showing sinus rhythm with ST segment elevation in leads aVR, V1 and widespread ST segment depression concerning for acute coronary syndrome

Video 2 Aortic root angiography using 6F Expo 145 Pig Catheter (Boston Scientific) showing a dissection flap in the proximal ascending aorta with extension to the arch. The flap prolapses into the left ventricle in diastole causing significant regurgitation of contrast material into the left ventricle

Video 1 Coronary angiography in the left anterior oblique projection using 5F EXPO FL5 Catheter (Boston Scientific). The catheter tip appears in the false lumen and dives towards the origin of the left main coronary artery

Video 3 Intraoperative transoesophageal echocardiography in mid-oesophageal, aortic valve long-axis view

aVR and V1, with widespread ST segment depression (figure 1). He underwent emergent coronary angiogram with initial diagnosis of acute coronary syndrome and impending cardiogenic shock. Angiogram revealed type-A aortic dissection extending

CrossMark

To cite: Ibrahim M, Rashid J, Kurtz SC, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-221399

Video 4 Intraoperative transoesophageal echocardiography with colour Doppler, in mid-oesophageal, aortic valve long-axis view demonstrating dissection of the proximal aorta with diastolic prolapse of a cylinder-shaped intimal flap into the left ventricle. The prolapsing flap prevents closure of the aortic valve leaflets causing severe aortic regurgitation

to the aortic arch with concomitant aortic regurgitation (AR) (videos 1 and 2). On prompt recognition of aortic dissection, he was transferred to the operating room for emergent surgical repair. Intraoperative transoesophageal echocardiography (TEE) showed circumferential dissection of the proximal aorta with diastolic prolapse of a cylinder-shaped intimal flap into the left ventricle causing severe AR (videos 3–5). The dissection was successfully repaired using 30 mm Hemashield Platinum double velour woven graft (MAQUET, New Jersey, USA). Aortic valve leaflets were resuspended successfully (videos 6 and 7, figure 2) and the coronaries were intact.

Postoperative course was unremarkable. The patient was discharged home on postoperative day 5. During outpatient follow-up at 1 week and 9 month intervals, he continued to be

Video 5 Intraoperative transoesophageal echocardiography in mid-oesophageal, aortic valve short-axis view demonstrating circumferential intimal dissection. In diastole, the dissected flap is obstructing the ostium of the left main coronary artery leading to the ECG changes seen

Video 6 Postrepair, transoesophageal echocardiography in mid-oesophageal, aortic valve short-axis view

stable. A small-sized, non-disabling stroke was reported at his second follow-up appointment. Interval CT angiography of the aorta at 9 months showed stable postoperative changes.

Video 7 Postrepair, transoesophageal echocardiography with colour Doppler in mid-oesophageal, aortic valve short-axis view, demonstrating normal coaptation of the aortic valve leaflets with trace residual aortic regurgitation

Figure 2 Intraoperative photograph of the aortic root and aortic valve (A) before repair and (B) after placement of the Hemashield Platinum graft and resuspension of the aortic valve leaflets

Learning points

- ▶ Acute Stanford type A aortic dissection is a life-threatening condition that necessitates prompt diagnosis and management. If untreated, the mortality rates have been reported as high as 60% in the first 24 hours and 90% within 3 months.¹
- ▶ In acute aortic dissection, the dissection flap is generally oriented along the long axis of the aorta. Circumferential intimal dissection of the ascending aorta is a rare variant. In this case, the intima undergoes circumferential tear and is separated from the aortic wall forming an intimal tube with a 'windsock' appearance.² The intimal flap may telescope in a retrograde or antegrade fashion causing what is described as 'intimo-intimal intussusception'.
- ▶ A retrograde intimo-intimal intussusception and prolapse of the aortic flap into the left ventricle is a fatal form of type-A aortic dissection. The prolapsing flap could suffocate the coronary ostia and compromise aortic valve leaflets closure. This results in severe AR, rapid haemodynamic collapse and ECG changes consistent with occlusion of the left main coronary artery as expressed in our case.³
- ▶ Early diagnosis and prompt operative intervention are essential for favourable clinical outcome. Unfortunately, a timely diagnosis of this type of aortic dissection is difficult since the presenting symptoms are frequently suggestive of acute coronary syndrome (ACS) and the patients are rushed to undergo coronary angiogram which potentially delays appropriate care. In our case, based on the ECG and haemodynamic findings, the impression was ACS involving the left main coronary artery with impending cardiogenic shock. All efforts were directed towards accomplishing early revascularisation which is essential for mortality benefit. However, the downstream work up revealed an aortic dissection as the primary problem instead of ACS which mandated an alternative management.
- ▶ High index of suspicion for acute aortic dissection in patients presenting with acute chest pain, combined with detailed physical exam and prompt utilisation of non-invasive diagnostic imaging like echocardiography, CT or magnetic resonance angiography (if applicable from haemodynamic standpoint) are key factors in diagnosing this type of acute aortic dissection and avoiding delay in appropriate care.

Contributors MI wrote the manuscript and edited all videos and images. JR took care of the patient during admission and edited the manuscript. SCK performed the coronary angiogram and reviewed the manuscript. SM reviewed and edited the manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Lijoi A, Scarano F, Canale C, *et al.* Circumferential dissection of the ascending aorta with intimal intussusception. Case report and review of the literature. *Tex Heart Inst J* 1994;21:166–9.
- 2 Yang EH, Kwon MH, Mahajan A, *et al.* Circumferential type A aortic dissection and intimal intussusception of the aorta causing severe aortic regurgitation and obstruction of the left main coronary artery. *Echocardiography* 2013;30:E81–4.
- 3 Lajevardi SS, Sian K, Ward M, *et al.* Circumferential intimal tear in type A aortic dissection with intimo-intimal intussusception into left ventricle and left main coronary artery occlusion. *J Thorac Cardiovasc Surg* 2012;144:e21–3.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow