Rose thorn injury

Katarina Kieselova, Felicidade Santiago, Martinha Henrique

Department of Dermatology,
Centro Hospitalar Leiria, Leiria,
Portugal

Correspondence to

Dr Katarina Kieselova,
katarinakieselova@gmail.com

Accepted 16 August 2017

DESCRIPTION

A 71-year-old man, with a history of arterial hypertension, presented with a 4-month history of four painful progressively enlarging nodules on the dorsal aspect of the right hand. The patient reported that he had been pricked by a rose on the fourth finger approximately 3 weeks before the appearance of the first lesion on the site of the injury. On physical examination, we observed four erythematous nodules, 10–15 mm in diameter, in linear distribution extending from the hand proximally (figure 1). Two nodules presented superficial erosions with purulent secretion. No systemic symptoms were detected. Histopathology revealed epidermal hyperkeratosis and suppurative small granulomas with polymorphous dermal infiltrate and periodic acid-Schiff staining identified rare fungal spores (figure 2). Culture from the biopsy specimen was negative. Clinical presentation and histology were consistent with sporotrichosis, so the patient was given itraconazole 200 mg daily. After 3 months of treatment, only a mild improvement was observed; therefore, the dose was escalated to 200 mg twice daily during following 4 months with complete resolution of 3 lesions. The

Figure 1 Four nodules on the dorsal aspect of the hand and forearm.

Figure 2 (A) Histology shows suppurative granulomas in dermis with overlying acanthotic epidermis and the areas of pseudoepitheliomatous hyperplasia (H&E, ×100). (B) Small granulomas composed of histiocytes, neutrophils and giant multinucleated cells are observed (H&E, ×100). (C) Oval fungal spores and yeast-like forms may be seen (periodic acid-Schiff, ×400).

fourth lesion was excised completely. There was no recurrence of the disease after 2-year follow-up.

Sporothrix schenckii, a saprophytic fungus, causes chronic infection, most frequently limited to skin

CrossMark

To cite: Kieselova K, Santiago F, Henrique M. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-221757

Learning points

- ▶ Sporotrichosis is a rare chronic granulomatous mycotic infection caused by a bimorphic fungus *Sporothrix schenckii*. The most common form of sporotrichosis is a lymphocutaneous infection, which is caused by traumatic inoculation from contaminated plants, soil or through zoonotic transmission.
- ▶ The diagnosis of sporotrichosis is based on clinical presentation, epidemiological context together with laboratory studies (direct examination, culture and tissue histopathology).¹
- ▶ Itraconazole 200 mg daily for at least 3–6 months is the treatment of choice for the lymphocutaneous disease. Non-responders should be given itraconazole 200 mg twice daily, terbinafine 500 mg twice daily or saturated solution of potassium iodide if itraconazole is not available. Possible adverse side effects of the prolonged systemic antifungal treatment (headache, gastrointestinal, dermatological manifestation) or prolonged potassium iodide administration (endocrine, gastrointestinal and renal) require careful clinical and laboratory monitoring.¹

and superficial lymphatics. Sporotrichosis presents as reddish papules at the site of inoculation, usually 3 weeks after the penetrating injury, with progression through the lymphatics forming multiple cutaneous nodules.¹ Systemic sporotrichosis is rare and occurs from the haematogenous spread from the primary inoculation site or from the pulmonary sporotrichosis in immunocompromised patients.²

Contributors All persons who meet authorship criteria are listed as authors. KK wrote the manuscript, reviewed by FS. KK performed the literature review. FS and MH made the diagnosis and performed the complete investigation and management of the patient. All authors approved the final version of manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Barros MB, de Almeida Paes R, Schubach AO. *Sporothrix schenckii* and sporotrichosis. *Clin Microbiol Rev* 2011;24:633–54.
- 2 Mahajan VK. Sporotrichosis: an overview and therapeutic options. *Dermatol Res Pract* 2014;2014:1–13.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow