Cysticercosis of the eyelid

Amar Pujari, Mandeep S Bajaj, Seema Sen, Vaishali Rakheja

Department of Ophthalmology,
Dr. Rajendra Prasad Centre for
Ophthalmic Sciences, AIIMS,
New Delhi, India

Correspondence to

Dr Amar Pujari,
dramarpujari@gmail.com

Accepted 23 August 2017

DESCRIPTION

A 26-year-old male patient complained of well-defined cystic swelling on the conjunctival aspect of his left lower eyelid for the past 2 months. His medical history was not significant. Visual acuity was 20/20 in each eye with unremarkable anterior and posterior segments, without any extraocular movement restriction. Examination of the left lower lid on its conjunctival aspect revealed a well-defined cystic lesion with soft to firm in consistency and it was completely preseptal (figure 1A). A complete excision biopsy was performed under local anaesthesia. Histopathology showed a multilayered cyst with minimal surrounding fluid around the invaginated scolex (figure 1B, Yellow arrow), four visible eccentric suckers (red arrow) and near central double layered hooklets (green arrow) suggestive of cysticercosis of the eyelid. Non-contrast enhanced CT of the brain and orbit did not reveal any additional foci of infection. The patient was prescribed oral prednisolone 1 mg/kg tapered over 4 weeks along with oral albendazole 15 mg/kg body weight for 4 weeks. At the end of 10 days, there was a healthy wound site (figure 1C) and at the end of 5 months, there was no recurrence/complications.

Figure 1 (A) A well-defined cystic lesion along the conjunctival aspect of the left lower eyelid. (B) Histopathology of the cyst showing a well-defined cyst having invaginated scolex (yellow arrow) with four eccentric suckers (red arrow) and a central double row of hooklets (green arrow). (C) At the end of 10 days, the wound is completely healed without much scarring of the conjunctiva.

Human cysticercosis is due to larval infestation by the cestode *Taenia solium*. The ocular involvement can be intraocular involving the vitreous or the subretinal space or as extra-ocular foci mainly involving the muscles.¹ The presentation in these cases can be as recurrent ocular surface inflammation, ptosis or self-limiting extraocular motility restriction. Involvement of the eyelid is rare, but these cases may also present as a subconjunctival mass or as spontaneous extrusion of the cyst. In the current case, the diagnosis of cysticercosis at first was unlikely as it was in an unusual location without any signs of inflammation. We kept a provisional diagnosis of the inclusion cyst before excision. As it was a cosmetic concern for the patient, the excision biopsy was performed with due consent. Postoperative histopathological examination proved vital in this case.

Learning points

- ▶ Periorbital involvement of the cysticercosis can have varied presentations. Under these circumstances, the final diagnosis is mainly achieved by a careful clinical examination, orbital imaging and monitored medical/surgical management.
- ▶ The differential diagnosis for orbital conditions includes many possibilities and can be a Pandora's box. Any form of tissue excision around the orbit is better subjected to a histopathological examination to ascertain the final pathology.

Contributors AP, MSB, SS and VR: have evaluated the case initially. AP: did excisional biopsy followed appropriate histopathological analysis. After analysing the educational value all authors together wrote the report.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCE

- 1 Pushker N, Bajaj MS, Betharia SM. Orbital and adnexal cysticercosis. *Clin Exp Ophthalmol* 2002;30:322–33.

CrossMark

To cite: Pujari A, Bajaj MS, Sen S, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-221823

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow