Sterile subdural empyema: an unusual presentation

Mohammed Al Salihi,^{1,2} Babikir Kheiri,^{1,2} Emad Abu Sitta,^{1,2} Abed Kanzy³

¹Internal Medicine Department, Hurley Medical Center, Flint, Michigan, USA

²Michigan state University

³Department of Internal Medicine, Hurley Medical Center, Flint, Michigan, USA

Correspondence to

Dr Emad Abu Sitta, emademad70@hotmail.com

Accepted 13 October 2017

DESCRIPTION

A 66-year-old man presented to the emergency department with a 6-week history of progressively worsening generalised weakness and gait instability. Physical examinations revealed a low-grade fever of 37.7°C. In addition, the patient was found to have very poor dentition and bilateral sensorineural hearing reduction with no other focal neurological deficit. Laboratory investigations showed a white cell count of $25.8 \times 10^9/L$ (Ref: $4-10.8 \times 10^9/L$), predominately neutrophils at 84% (Ref: 36%–75%). A CT of the head was negative for any acute processes and a subsequent MRI of the head with and without contrast was consistent with bilateral frontal subdural empyema (SDE) and ventriculitis (figure 1). He was started initially on vancomycin, ceftriaxone and metronidazole and he underwent bilateral burr hole biopsy and drainage of a thick, creamy white fluid within 2 days of starting the antibiotics. Pathological findings were consistent with an abscess; however, the microbiological culture failed to grow any organisms. The patient was continued on ceftriaxone and metronidazole for a total of 5 weeks. He made very well recovery with no residual neurological deficits.

SDE is a rare, life-threatening, purulent collection of material between the outer dura mater and inner arachnoid mater.¹ It accounts for 20% of all cases of intracranial abscesses and usually occurs following sinusitis, head trauma or surgery,

Learning points

- ▶ Subdural empyema (SDE) is a rare, life-threatening neurological emergency and if left untreated can eventually lead to death.
- ▶ Early recognition and rapid antibiotics and surgical intervention can reduce the morbidity and mortality of SDE.

metastatic bacteraemia or pre-existing subdural haematoma.¹ In this case, it was attributed to poor dentition, which accounts for only 0.7% of SDE cases.² Although operative cultures can identify the causative organisms, 7%–52% of cases yield no growth, which is largely attributed to the prior use of antibiotics or improper use of anaerobic culture techniques.³

SDE is a true neuroglial emergency³ and, if untreated, can lead to permanent neurological deficits such as seizures, cerebral venous thrombosis, hydrocephalus, cerebral swelling, coma and eventually death.¹ Diagnosis is made with radiographic imaging including CT scan and, more sensitive, MRI.^{1,3} Treatment requires a prolonged course of antibiotics and immediate surgical evacuation with either a burr hole or craniotomy.³ These measures have led to reduced morbidity and mortality.¹

Contributors MAS, resident on the case, wrote the manuscript and reviewed the literature. BK, resident on the case, reviewed the manuscript and reviewed the literature. AK, primary care of the patient, reviewed the manuscript. EAS, attending, reviewed the manuscript and reviewed the literature.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

Figure 1 MRI head. (A) Fast spoiled gradient recalled 3D sequence with contrast showing bilateral frontal subdural fluid collections measuring up to 1.1 cm on the left and 1 cm on the right (arrows). (B) Axial T1 weighted with contrast showing signal abnormalities within the occipital horns of both lateral ventricles consistent with ventriculitis (arrows).

REFERENCES

- 1 Agrawal A, Timothy J, Pandit L, et al. A review of subdural empyema and its management. *Infectious Diseases in Clinical Practice* 2007;15:149–53.
- 2 Nathoo N, Nadvi SS, van Dellen JR, et al. Intracranial subdural empyemas in the era of computed tomography: a review of 699 cases. *Neurosurgery* 1999;44:529–35.
- 3 Osborn MK, Steinberg JP. Subdural empyema and other suppurative complications of paranasal sinusitis. *Lancet Infect Dis* 2007;7:62–7.

CrossMark

To cite: Al Salihi M, Kheiri B, Abu Sitta E, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-222053

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow