Unilateral erythema nodosum: atypical presentation in paediatrics

Francisca Calheiros-Trigo, Sofia Martins

Department of Pediatrics,
Hospital de Braga, Braga,
Portugal

Correspondence to
Dr Francisca Calheiros-Trigo,
trigo.francisca@gmail.com

Accepted 10 November 2017

DESCRIPTION

A previously healthy 12-year-old boy was observed in the emergency department due to pain and erythema in the left shin for the past 2 weeks. He was discharged with flucloxacilin for cellulitis. One week later, he returned with oedema and erythema of the left shin, with palpable nodules and purple discolouration (figure 1). The right leg was normal. Oropharyngeal hyperaemia was observed.

Complementary study revealed erythrocyte sedimentation rate of 12 mm/hour and antistreptolysin O titre was 985 UI/mL (normal range 0–408 UI/mL). C reactive protein was <2.90 mg/dL; autoimmunity study, infectious serologies (hepatitis B virus and Epstein-Barr virus, EBV, *Mycoplasma pneumoniae* and *Salmonella* spp.) rapid strep test and Mantoux test were negative. Ultrasound showed subcutaneous oedema. Biopsy revealed septal panniculitis, compatible with erythema nodosum (figure 2). He was discharged.

Four weeks later, there was resolution of nodules and oedema of left shin, which showed only minor discolouration, without atrophy or scarring (figure 3).

Erythema nodosum is the most common presentation of panniculitis.^{1 2} This disease is rare among children, and its peak incidence occurs in the third decade.^{1–3} We found no descriptions of unilateral erythema nodosum in children. Different stimuli cause inflammation of subcutaneous fat.^{1–3} Usually, no underlying

Figure 2 Incisional biopsy. H&E ×40. Septal panniculitis.

Figure 3 Re-evaluation 4 weeks after discharge.

Figure 1 Erythema, oedema and nodules in the left shin.

cause is identifiable, but systemic inflammatory diseases, infectious diseases, neoplasm and drug reactions should be excluded.¹ The most common provoking insult in children is β -haemolytic *Streptococcus*.³ Diagnosis is clinical: bilateral erythematous nodules, frequently in the shins, that evolve in the course of 4–8 weeks with complete resolution.¹ When the diagnostic criteria are not fulfilled, biopsy must be performed to clarify doubtful cases.^{1 3}

Learning points

- ▶ Erythema nodosum is diagnosed based on clinical criteria: bilateral erythematous nodules, most frequently in the pretibial region.
- ▶ When diagnostic criteria are not fulfilled (eg, unilateral lesions), incisional biopsy must be performed for diagnosis.
- ▶ Unilateral erythema nodosum is rare, but one must be aware of this entity.

CrossMark

To cite: Calheiros-Trigo F, Martins S. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-222441

Contributors Report written by FCT and overseen by SM.

Competing interests None declared.

Patient consent Guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Schwartz RA, Nervi SJ. Erythema nodosum: a sign of systemic disease. *Am Fam Physician* 2007;75:695–700.
- 2 Litwin L, Machura E. The etiology and clinical manifestation of erythema nodosum in hospitalized children - analysis of 12 cases. Preliminary report. *Dev Period Med* 2014;18:506–12.
- 3 Kakourou T, Drosatou P, Psychou F, et al. Erythema nodosum in children: a prospective study. *J Am Acad Dermatol* 2001;44:17–21.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow