Complication of deep brain stimulation for Parkinson's disease

Catarina Costa,¹ Felisbela Gomes,¹ Joaquim Monteiro,² Luís Bento³

¹Medicina 2.1, Centro Hospitalar de Lisboa Central EPE, Lisboa, Portugal

²Servico de Neurocirurgia, Centro Hospitalar de Lisboa Central EPE, Lisboa, Portugal

³Unidade de Urgencia Medica, Hospital de Sao Jose, Lisboa, Portugal

Correspondence to

Dr Catarina Costa,
costa.catarina@gmail.com

Accepted 19 November 2017

DESCRIPTION

A 58-year-old man with advanced Parkinson disease diagnosed 13 years before, was admitted for surgical placing of continuous bilateral high-frequency stimulation of the subthalamic nucleus.

The procedure was uneventful (lead used Medtronic DBS 3387; antibiotic prophylaxis: cefazolin 2g at time of anaesthetic induction and 1g every 8 hours in the first 24 hours) with a post-operative control CT scan showing good placement of the electrodes and no abnormalities (figure 1). The stimulation was switched on 72 hours after the procedure and within 24 hours the patient presented two epileptic seizures controlled with diazepam and stimulation was switched off. The head CT showed intra-axial hypodensity more prominent in the frontal area with a gaseous collection and oedema surrounding the right electrode pathway (figure 2). At the time there was no fever, no increase in inflammatory markers and a normal cell count in the lumbar puncture. However, antibiotic therapy (ceftriaxone and metronidazole) and phenytoin were started.

The depressed mental status and maintenance of epileptic crisis motivated sedation, orotracheal intubation and admission to the intensive care unit. Vancomycin and levetiracetam were added to the already ongoing therapy. An electroencephalography showed diffuse slowing down without paroxysmic activity and

Figure 2 Head CT scan showing intra-axial hypodensity (more prominent in the frontal area) with a gaseous collection and oedema surrounding the right electrode pathway.

a CT angiography excluded venous and arterial thrombosis. The electrodes were not removed. All cultural exams were negative and sequential head CT scans showed a reduction in the hypodensity area with resorption of the gaseous collection and diminished oedema (figure 3).

Figure 1 Postoperative head CT scan showing good placement of the electrodes.

Figure 3 Head CT scan showing a reduction in the hypodensity area with resorption of the gaseous collection and diminished oedema.

CrossMark

To cite: Costa C, Gomes F, Monteiro J, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-223264

Learning points

- Deep brain stimulation is a routine surgical treatment used in selected cases of Parkinson's disease.
- Complications after electrode implantation are described, and in this case, we admitted surrounding oedema but we could not rule out infection.^{1–3} The unilateral electrode surrounding oedema is a rare complication but excludes an allergic reaction to the material and a complication of the stimulation itself.
- The decision not to remove the electrodes turned out to be beneficial as the patient profited from the stimulation after this episode.

There was a progressive recovery of the neurological state without sequelae allowing transfer to the neurology ward and discharge from the hospital.

Contributors CC was responsible for the conduction, reporting and conception of the article. FG was responsible for the acquisition of data. JM and LB were involved in the planning and revision.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Schoen NB, Jermakowicz WJ, Luca CC, *et al.* Acute symptomatic peri-lead edema 33 hours after deep brain stimulation surgery: a case report. *J Med Case Rep* 2017;11:103.
- 2 Pouratian N, Reames DL, Frysinger R, *et al.* Comprehensive analysis of risk factors for seizures after deep brain stimulation surgery. Clinical article. *J Neurosurg* 2011;115:310–5.
- 3 Fily F, Haegelen C, Tattevin P, *et al.* Deep brain stimulation hardware-related infections: a report of 12 cases and review of the literature. *Clin Infect Dis* 2011;52:1020–3.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow