A rare cause of neonatal persistent jaundice

Gracinda Nogueira Oliveira,¹ Isabel Dinis,² Maria José Noruegas,³ Alice Mirante²

¹Department of Paediatrics, Hospital Pediátrico de Coimbra, Coimbra, Portugal

²Department of Paediatric Endocrinology, Diabetes and Growth, Hospital Pediátrico de Coimbra, Coimbra, Portugal

³Department of Radiology, Hospital Pediátrico de Coimbra, Coimbra, Portugal

Correspondence to

Dr Gracinda Nogueira Oliveira, gracenoliveira@gmail.com

Accepted 9 December 2017

DESCRIPTION

A 22-year-old gravida 2, para 1 (G2P1) woman with immunoglobulin anti-D prophylaxis, insulin-treated gestational diabetes and first-trimester cytomegalovirus (CMV) infection vaginally delivered a 39-week boy weighing 3720 g (90th centile) and with Apgar scores of 8 and 10 at 1 and 5 min. Prenatal ultrasonographic assessment throughout gestation was normal. Nursery stay was uneventful. He was discharged on day 2, with a normal examination, except for the appearance of jaundice, with a transcutaneous bilirubin of 248 $\mu\text{mol/L}$ (cut-off 250 $\mu\text{mol/L}$), not meeting the criteria for phototherapy. A follow-up clinic on day 4, arranged for bilirubin measurement and CMV testing, surprisingly revealed poor general appearance, lethargy, very icteric skin and a minor weight loss (9% of birth weight). Both liver and splenic edges were palpable. Vital signs were normal. Blood routine showed haemoglobin of 19 g/dL, haematocrit of 58%, white blood cells $11.6 \times 10^9/\text{L}$ and platelet count $128 \times 10^9/\text{L}$. Biochemistry revealed total serum bilirubin of 694 $\mu\text{mol/L}$ (cut-off 350 $\mu\text{mol/L}$), indirect bilirubin of 681 $\mu\text{mol/L}$, and a normal glucose, urea, creatinine and coagulation study. He was admitted for intensive phototherapy, and his total serum bilirubin level decreased to

289 $\mu\text{mol/L}$ in 9 hours, but rebound hyperbilirubinaemia followed whenever attempts to discontinue were made. He remained icteric, hypotonic, required enteral tube feeding and had a persistent hyponatraemia treated with oral sodium chloride. On day 9 cranial ultrasound (US) was normal, but abdominal US uncovered enlarged adrenal glands, with a hypoechoic heterogeneous mass, measuring $32 \times 21 \times 17 \text{ mm}$ on the right and $33 \times 23 \times 17 \text{ mm}$ on the left, suggesting a bilateral adrenal haemorrhage (AH). CMV viruria came negative, and subsequent studies showed increased thyroid-stimulating hormone (TSH) and normal free thyroxine 4 levels, leading to the initiation of levothyroxine treatment on day 14. Transfer to a tertiary paediatric hospital was arranged on day 28. Abdominal Doppler US on day 30 confirmed the bilateral AH (figure 1A). The complete hormone evaluation showed raised adrenocorticotrophic hormone (ACTH), prolactin, aldosterone and renin, and low cortisol level, which, together with an elevated TSH, disclosed hypothyroidism secondary to adrenal insufficiency (AI). Under hydrocortisone treatment, started on day 39, hypotonia resolved and feeding improved with an adequate weight gain. Neurological and metabolic evaluations were normal. Thyroid parameters normalised at 3 months and levothyroxine was suspended. At 6 months fludrocortisone was added to control the inappropriately activated renin-angiotensin-aldosterone system. Two-month and 6-month follow-up US revealed an ongoing and significant regression of both adrenal masses in size and echogenicity (figure 1B,C). Presently, at 22 months, he remains asymptomatic but continues on corticosteroid replacement therapy for hormonal control. The most recent US, at 12 months, showed an even smaller bilateral haemorrhage with a maximum diameter of 6 mm.

AH is a rare neonatal event, most often unilateral and located on the right side (70%) and infrequently bilateral (5%–10%).^{1,2} Predisposing factors include difficult delivery, maternal diabetes, large for gestational age infants, birth asphyxia or trauma, septicaemia and underlying tumour.^{2,3} The aetiology of our patient's bilateral AH is unknown. AH clinical manifestations are non-specific and vary according to the degree of haemorrhage and subsequent adrenal cortex compromise.² Neonatal jaundice and flank mass are the most common presentations. The main differential diagnosis of AH is the cystic form of neuroblastoma; thus, vanillylmandelic acid urinary levels should be assessed if colour Doppler reveals blood flow over the mass.^{2,3} Doppler US is the diagnostic tool of choice for initial and follow-up evaluation of neonates with suspected AH.^{1,2,3} Nevertheless, the authors consider that inconclusive US findings, progression in lesion size

Figure 1 (A) One-month ultrasound (US): hypoechoic and heterogeneous ovoid mass with no blood flow at colour Doppler and surrounding calcification compatible with haemorrhage. (B) Two-month US shows at least a 50% bilateral decrease in the cystic mass area and echogenicity. (C) Six-month US shows further bilateral decrease in size and echogenicity. L, left adrenal gland; R, right adrenal gland.

CrossMark

To cite: Oliveira GN, Dinis I, Noruegas MJ, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-223306

Images in...

or unusual clinical courses justify the use of MRI to confirm the presence of haemorrhage and determine if blood is the sole component of the haematoma. On its turn, CT involves ionising radiation exposure, and therefore we consider that it should be reserved primarily for emergent cases. Although there is no specific US sign for AH, demonstrating the absence of vascular supply and a decrease in lesion size can be considered sufficient for AH diagnosis and avoid further investigation in otherwise normally formed term newborns with AI (as in our case).¹

Both serum cortisol and ACTH should be measured in newborns with AH to exclude primary AI.² Hyponatraemia in

our patient was most likely the result of hypervolaemia due to an increased antidiuretic hormone (ADH) secretion elicited by cortisol deficiency. AI in a newborn with bilateral AH requires supplementation with both glucocorticoids and mineralocorticoids.² A close follow-up using serial US is required, and AH usually resolves between 3 weeks and 6 months.^{2 3} However, a longer period of time is possible in accordance with our case.

Contributors GNO conceptualised the case report, did a literature review, integrated the data and wrote the manuscript. ID and MJN contributed to the conception, drafting and critical revision of the article. AM closely supervised the management of the patient and the writing of the case report.

Competing interests None declared.

Patient consent Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Katar S, Öztürkmen-Akay H, Devecioğlu C, *et al.* A rare cause of hyperbilirubinemia in a newborn: bilateral adrenal hematoma. *Turk J Pediatr* 2008;50:485–7.
- 2 Qureshi UA, Ahmad N, Rasool A, *et al.* Neonatal adrenal hemorrhage presenting as late onset neonatal jaundice. *J Indian Assoc Pediatr Surg* 2009;14:221–3.
- 3 Mutlu M, Karagüzel G, Aslan Y, *et al.* Adrenal hemorrhage in newborns: a retrospective study. *World J Pediatr* 2011;7:355–7.

Learning points

- Bilateral adrenal haemorrhage (AH) is a rare neonatal event that should be recognised in the presence of persistent unexplained jaundice and predisposing factors.
- Abdominal Doppler ultrasound is the examination of choice in neonates with suspected AH, allowing organ injury assessment and exclusion of other abnormalities at diagnosis and a non-invasive monitoring at follow-up.
- AH is usually self-limiting, but rare bilateral cases may cause adrenal insufficiency and require corticosteroid therapy.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>. BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow