Beware of air

Nadia, NK van Rijssel, Claudia, C Savelkoul, David H T Tjan

Department of Intensive Care
Unit, Gelderse Vallei Hospital,
Ede, The Netherlands

Correspondence to

Drs David H T Tjan, tjand@zgv.nl

NK contributed equally,
C contributed equally.

Accepted 1 November 2017

DESCRIPTION

We present three patients, treated at our intensive care unit (ICU), with radiological signs of free gas in different tissues.

Patient A, a 67-year-old man with a history of *colitis ulcerosa*, was admitted with acute right upper quadrant pain and fever. An abdominal X-ray (figure 1) showed free gas under the diaphragm and Rigler's sign (indicating the presence of gas on both sides of the bowel wall).¹ The CT scan revealed a toxic megacolon with signs of a pneumoperitoneum (figure 2) caused by a colon perforation.

Figure 1 Free gas in the abdominal cavity and Rigler's sign (indicated by white arrows).

Figure 2 A toxic megacolon with signs of a pneumoperitoneum (indicated by red arrows).

Patient B, a healthy 62-year-old man, presented with malaise and substernal chest pain. During physical examination, a retropharyngeal abscess was found. The CT scan revealed an extensive pneumomediastinum with gas penetrating into the nuchal region (figure 3). Both patients showed signs of sepsis with tachypnoea, tachycardia and fever.

Patient C, a 78-year-old woman with Addison's disease, complained of subacute fever and back pain. Her CT scan revealed a pneumomyositis and pneumovertebralis (respectively intramuscular and intravertebral gas) caused by a perforated diverticulitis with a fistula (figure 4). However, her vital signs were stable. Suppression of septic shock due to a maintenance dosage of prednisolone was therefore suspected.

All patients were immediately treated with broad-spectrum antibiotics and underwent subsequent surgical exploration. Patients A and C had a pneumoperitoneum, indicating free gas trapped within the peritoneal cavity but outside

Figure 3 An extensive pneumomediastinum and gas penetrating into the nuchal region (indicated by red arrows).

CrossMark

To cite: van Rijssel N,NK,
Savelkoul C,C, Tjan DHT. *BMJ*
Case Rep Published Online
First: [please include Day
Month Year]. doi:10.1136/
bcr-2017-223081

Figure 4 Pneumomyelitis and pneumovertebralis (indicated by red arrows).

the lumen of the bowel. Patient B had a pneumomediastinum due to mediastinitis. Afterwards, they were admitted to the ICU for postoperative care. Patients A and B both went into severe septic shock. Against all odds, patient C recovered without any clinical sequelae.

A complete summary of differential diagnoses regarding free gas can be found in [table 1](#). Up to 49% of cases of a pneumoperitoneum may not be detected using conventional radiography. Therefore, it is recommended to make a CT scan, which can correctly identify the site of perforation in 86% of the cases.² After taking microbiological cultures, broad-spectrum antibiotics should be started immediately if sepsis is suspected. However, there may also be causes of free gas for which antibiotics are not the direct course of treatment. Surgical exploration is mandatory if sepsis source control is required.

Radiological images showing free gas are alarming as this may indicate a possible fulminant course of disease. In these cases, an initial aggressive approach is justified. Striking images of patients presenting with free gas, however, are not always concordant with a rapid disease progression.

Table 1 Differential diagnosis of free gas

Tract	Origin
Airway	Drugs (eg, cocaine)
	Pneumothorax
	▶ asthma/COPD
	▶ idiopathic
	▶ iatrogenic (eg, mechanical ventilation)
Gastrointestinal	▶ trauma
	Retropharyngeal abscess
	Trauma of the upper respiratory tract
	Abscess
	Appendicitis
	Bacterial peritonitis
	Diverticulitis
	Foreign bodies
	(Incarcerated) hernia
	▶ abdominal wall hernia
	▶ hiatus hernia
	Iatrogenic
	▶ drugs (eg, NSAIDs)
	▶ endoscopy
	▶ paracentesis
	▶ peritoneal dialysis
	▶ postirradiation
	▶ surgery
	Inflammatory bowel disease
	Ischaemic bowel
	Malignancy
	▶ primary
	▶ metastatic
	Necrotising enterocolitis
	Peptic ulcer disease
	Reflux esophagitis
	Trauma
	Volvulus
	▶ after prior abdominal surgery
	▶ congenital
Soft tissue	Gas from other tract
	▶ airway
	▶ gastrointestinal
	Infectious
	▶ fasciitis necroticans
	▶ toxic shock syndrome
	Iatrogenic
Vascular	Trauma
	Diving
	Iatrogenic

COPD, chronic obstructive pulmonary disease; NSAIDs, non-steroidal anti-inflammatory drugs.

Therefore, the interpretation of clinical findings and radiological images should always be combined for appropriate decision-making.

In conclusion, due to the possible severe course of disease, every patient with free gas should be approached with caution. Depending on whether there is a suspicion of sepsis,

Learning points

- ▶ Imaging showing free gas warrants a high index of suspicion; however, striking images are not always concordant with the course of the disease.
- ▶ Free gas has different anatomical origins and not all causes are life threatening.
- ▶ Sepsis as a cause requires antibiotic treatment and usually surgical source control.

immediate source control through explorative surgery and postoperative ICU care should be considered.

Contributors NvR participated in writing the article (and revision) and received informed consent from the patient. CS participated in writing the article (and revision). NvR, CS and DHTT were closely involved in patients' stay at ICU. DHTT critically revised the article. NvR and CS are guarantors and contributed equally.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Lloyd-Jones G. Abdominal X-ray – Abnormal bowel gas pattern [internet]. *Radiology Masterclass* 2007. http://www.radiologymasterclass.co.uk/tutorials/abdo/abdomen_x-ray_abnormalities/pathology_bowel_gas_perforation (cited 04 October 2017).
- 2 Coppolino F, Gatta G, Di Grezia G, *et al.* Gastrointestinal perforation: ultrasonographic diagnosis. *Crit Ultrasound J* 2013;5(Suppl 1):S4.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow