

Brain abscess in a patient with chronic sinusitis

Marta de Sousa,¹ Alice Lança,² Carolina Sepúlveda,¹ Edgar Pereira¹

¹Internal Medicine, Centro Hospitalar do Médio Tejo EPE Unidade de Torres Novas, Torres Novas, Portugal

²Nephrology, Centro Hospitalar do Médio Tejo EPE Unidade de Torres Novas, Torres Novas, Portugal

Correspondence to

Dr Alice Lança,
alancelancabaptista@gmail.com

Accepted 20 December 2017

DESCRIPTION

A middle-aged patient with a previous history of chronic sinusitis and arterial hypertension was admitted because of altered mental status. Two days earlier, symptoms such as fever, chills, nasal obstruction and headache slowly developed. On examination, the patient was febrile and with stiff neck. The remainder of the physical examination was normal. CT scan (figure 1A,B) and T1-MRI (figure 2A–C) were both performed; besides signs of sinusitis, a 2 cm right peritrial expansive lesion surrounded by vasogenic oedema was found being compatible with brain abscess. A thorough investigation was done to exclude infectious endocarditis, including echocardiography and blood cultures which were all negative. Viral serologies, including HIV, were negative. Based on the brain lesion characteristics (size and peculiar location), a conservative approach with broad-spectrum antibiotics (ceftriaxone, metronidazole and vancomycin) was initially suggested by Neurosurgery. However, 20 days after, the patient presented with neurological deterioration. A CT scan was repeated showing ventriculomegaly and hydrocephalus which required extraventricular drain placement. All blood cultures remained negative. Unfortunately, despite all efforts, the patient died 2 days later.

Intracranial abscess is an uncommon, serious and life-threatening infection with poor outcome.¹ The classic clinical triad of fever, headache and focal neurological deficits should raise the suspicion of brain abscess despite occurring in only about 2–34% of patients.¹ Aetiology can often be identified; however, in about 15% of cases, no source can be found. Occasionally, brain abscess results from contiguous infections such as sinusitis.¹ Based on the patient's clinical presentation, sinusitis was the most probable source of infection.




Figure 2 (A) MRI coronal plane demonstrating peritrial abscess and right posterior lateral ventriculitis. (B) MRI axial plane demonstrating peritrial abscess and right posterior lateral ventriculitis. (C) Another perspective of the peritrial abscess and right posterior lateral ventriculitis in MRI sagittal plane. (A–C) MRI demonstrating peritrial abscess and right posterior lateral ventriculitis.

According to the literature, in pyogenic abscess <2.5 cm, the first-line treatment is antimicrobial therapy; for those bigger than 2.5 cm, surgical intervention (adequate drainage) is recommended.² However, this decision should be individualised based on the evidence of growing abscess while on antibiotics or no change in size at 2–3 weeks.^{2,3} In this case, ventricular drainage combined with intravenous and/or intrathecal antibiotics should be immediately performed.³ Mortality rate correlates directly with the disease progression and patients' neurological status on admission.³




Figure 1 (A) CT scan coronal plane showing brain abscess surrounded by vasogenic oedema. (B) CT scan sagittal plane showing brain abscess surrounded by vasogenic oedema. (A,B) CT scan showing brain abscess surrounded by vasogenic oedema.



CrossMark

To cite: de Sousa M, Lança A, Sepúlveda C, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-223266

Learning points

- ▶ High clinical suspicion is necessary for early recognition and prompt treatment.
- ▶ Especially in immunocompetent patients, infection from contiguous structures must be considered.
- ▶ Despite the advent of newer antibiotics and surgical strategies, the optimal clinical approach remains controversial.

Contributors MdS, AL and CS made the diagnosis and wrote the manuscript. EP reviewed the clinical case and the manuscript.

Competing interests None declared.

Patient consent Detail has been removed from this case description/these case descriptions to ensure anonymity. The editors and reviewers have seen the detailed information available and are satisfied that the information backs up the case the authors are making.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Patel K, Clifford DB. Bacterial brain abscess. *Neurohospitalist* 2014;4:196–204.
- 2 Arlotti M, Grossi P, Pea F, *et al.* Consensus document on controversial issues for the treatment of infections of the central nervous system: bacterial brain abscesses. *Int J Infect Dis* 2010;14:S79–92.
- 3 Alvis Miranda H, Castellar-Leones SM, Elzain MA, *et al.* Brain abscess: current management. *J Neurosci Rural Pract* 2013;4:S67–81.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow