A rare involvement of epitrochlear lymph nodes in mantle cell lymphoma

Mohan Kumar, Gaurav Prakash, Kundan Mishra, Savita Kumari

Department of Internal Medicine, Post Graduate Institute of Medical Education and Research, Chandigarh, India

Correspondence to
Mohan Kumar,
monu8501@gmail.com

Accepted 9 April 2018

DESCRIPTION

A 60-year-old man presented with bilateral groin swelling for 1 year. Sequentially, he developed similar swellings in the cervical and axillary regions. He noted multiple nodular swellings arranged on the medial aspect of both arms in a linear fashion. At this moment, he reported to our hospital. He had generalised lymphadenopathy and the most characteristic finding was enlarged, bilateral, firm, epitrochlear lymph nodes ([figure 1](#)). No finding of local temperature rise, redness or infection was found. He also had hepatosplenomegaly. An excisional biopsy revealed mantle cell lymphoma. Bone marrow aspiration done for staging showed infiltration by mantle cells.

Epitrochlear lymph nodes are present on the medial aspect of the arm, about 1–2 cm above the elbow joint anterior to the medial intermuscular septum between the biceps and triceps.¹ Their enlargement is usually a part of generalised lymphadenopathy and sometimes because of pathologies in the hand and forearm. One study from the University of Edinburgh showed no palpable lymph nodes in 140 healthy individuals but nodes were palpable in 27% of 184 patients who were diagnosed with disease causing generalised lymphadenopathy.² Enlargement of epitrochlear lymph nodes is almost

always pathological. Isolated enlargement of epitrochlear lymph nodes has also been reported.³ They can be enlarged in infectious conditions like HIV, leishmaniasis, syphilis, leprosy, cat scratch disease, tuberculosis, filariasis and inflammatory conditions like rheumatoid arthritis when the disease is active in the hands. Their enlargement is much more common in chronic lymphoproliferative disorders than in aggressive lymphomas.

Learning points

- ▶ This case highlights the importance of clinical examination of the sites we usually miss during our routine clinics.
- ▶ It is an easily accessible site for biopsy.
- ▶ Although a less common clinical finding, enlarged epitrochlear lymph nodes can provide clues for evaluation of generalised lymphadenopathy and febrile illnesses.

Contributors MKH, GP, KM and SK were involved in the management of the patient. MKH and KM wrote the manuscript which was vetted by all.

Funding This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Fujiwara M, Suzuki A, Mizukami T, *et al*. Mid-arm lymph nodes dissection for melanoma. *J Plast Reconstr Aesthet Surg* 2010;63:1561–4.
- 2 Selby CD, Marcus HS, Toghill PJ. Enlarged epitrochlear lymph nodes: an old physical sign revisited. *J R Coll Physicians Lond* 1992;26:159–61.
- 3 Yardimci VH, Yardimci AH. An unusual first manifestation of hodgkin lymphoma: epitrochlear lymph node involvement-a case report and brief review of literature. *J Investig Med High Impact Case Rep* 2017;5:232470961770670.

Figure 1 Clinical photograph showing bilateral epitrochlear lymphadenopathy.

To cite: Kumar M, Prakash G, Mishra K, *et al*. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-224665

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow