Unique presentation of a ruptured Meckel's diverticulum after blunt trauma

Derek Lim, Kevin Bain, Sampath Kumar

Department of Surgery, NYU Langone Hospital - Brooklyn, Brooklyn, New York, USA

Correspondence to Dr Kevin Bain, kevin.bain@nyumc.org

Accepted 1 May 2018

DESCRIPTION

A 25-year-old man with no significant medical or surgical history presented with a complaint of thick, dark drainage from a periumbilical wound for 1 month (figure 1). He reported first noticing the drainage several days after bumping into the corner of a machine at work. He was seen by his primary care physician and prescribed oral antibiotics for persistent drainage and erythema; however, his symptoms failed to improve.

On presentation to the emergency department, the patient was afebrile and haemodynamically stable. Lab work was within normal limits. An abdominal CT scan demonstrated a loculated periumbilical fluid collection, with extension intraperitoneally into a portion of thickened small bowel with suspicion of an enterocutaneous fistula (figure 2).

The decision was made to take the patient to the operating theatre for further examination. Intraoperatively, the presence of a loop of small bowel densely adherent to the umbilicus was identified. After careful dissection, there appeared to be a perforated diverticulum located in proximity to the ileocaecal junction. The edges of the surrounding bowel were noted to be necrotic, and the decision was made to resect the segment. A small bowel resection was performed with primary anastomosis. The remainder of the patient's hospital course was uncomplicated. Surgical pathology confirmed a 2.5 cm Meckel's diverticulum: small intestine with a single diverticulum showing extensive mural gastric and pancreatic tissue. Histology demonstrated normal mucosa without evidence of dysplasia or

Figure 2 Loculated periumbilical fluid collection approximately 3.8×2.2×3.4 cm. An intraperitoneal portion of the fluid collection is contiguous with a thickened small bowel loop, which is suspicious for an enterocutaneous fistula.

autodigestion. This suggests diverticular rupture secondary to mechanical force.

Meckel's diverticulum is an outpouching of the small intestine that results from incomplete closure of the vitelline duct. It is most commonly located 40–60 cm from the ileocaecal valve, and occurs in 2%–3% of the population.^{1 2} In most instances Meckel's diverticulum is benign, asymptomatic and often diagnosed incidentally during laparoscopy/laparotomy. Meckel's diverticulum may contain gastric or pancreatic mucosa, which can become symptomatic. Most commonly, patients with

Figure 1 Chronic umbilical wound sustained after blunt abdominal trauma, with drainage of a thick, dark fluid for 1 month.

Learning points

- ▶ Meckel's diverticulum is often asymptomatic and discovered incidentally.
- ▶ The most common presenting symptom is gastrointestinal bleeding, and this is usually secondary to gastric mucosa present within the diverticulum, resulting in an acid-induced ulcer adjacent to the diverticulum.²
- ▶ Traumatic small bowel injuries are most common after blunt traumas (ie, motor vehicle accidents/seat belts, contact sports), and result in an increase in intra-abdominal pressure, which can lead to perforation in susceptible regions such as in the diverticulum or the terminal ileum.^{1 3}
- ▶ The mainstay of treatment in symptomatic and traumatic Meckel's diverticulum is diverticulectomy, with or without bowel resection.
- ▶ Resection of asymptomatic diverticulum remains controversial, and the risks/benefits and overall health of the patient must be considered.²

To cite: Lim D, Bain K, Kumar S. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-224751

ectopic gastric mucosa form adjacent ulcers, which present as a painless lower gastrointestinal bleed.²

Traumatic rupture of a Meckel's diverticulum is a rare occurrence. Previous cases have described rupture secondary to blunt injury from motor vehicle accidents and contact sports (soccer, softball). The force of impact during these injuries was significant enough to cause acute rupture of the diverticulum as well as arterial avulsion. These patients presented with acute signs of peritonitis, including guarding, rigidity and haemodynamic instability.^{1,3} A proposed aetiology for this rupture to occur is due to the sudden increase in intra-abdominal pressure. As compression of organs occurs against the impacting object and vertebrae, increased intra-abdominal pressure may cause perforation in susceptible areas such as the ileum, rectum and blind loops.³

These acute physical exam findings were absent in our patient. This is the first case reporting a blunt traumatic rupture of a Meckel's diverticulum that presented as a chronic enterocutaneous fistula. It emphasises the importance of obtaining a thorough history and physical exam, along with proper imaging, to guide the formulation of an accurate diagnosis and treatment plan.

Contributors DL, KB: research, writing, editing, formatting. DL, KB, SK: final editing.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Kazemi K, Jalaieian H, Fattahi MR, *et al.* Ruptured Meckel's Mesodiverticulum and Meckel's Diverticulum following blunt abdominal trauma. *Medical Principles and Practice* 2007.
- 2 Townsend Jr C, Beauchamp DR, Evers MB, *et al.* *Sabiston textbook of surgery: the biological basis of modern surgical practice*. 1. 20th Edn. Philadelphia, PA: Elsevier, 2017.
- 3 Tummers WS, Van Der Vorst JR, Swank DJ. Traumatic Rupture of a Meckel's Diverticulum due to blunt abdominal trauma in a soccer game: a case report. *International Journal of Surgery Case Report* 2015.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow