Pineal gland apoplexy mimicking as migraine-like headache

Davala Krishna Karthik,¹ Vikram Khardenavis,² Sharvari Kulkarni,³ Anirudda Deshpande⁴

¹Department of Radiology, Surya Diagnostics, Warangal, Telangana, India

²Department of Medicine, Miraj Medical College, Miraj, Maharashtra, India

³Department of Medicine, Maharashtra Institute of Medical Education and Research, Pune, Maharashtra, India

⁴Department of Neurology, Vinayaka Neuro Multispecialty Clinic, Warangal, India

Correspondence to

Dr Anirudda Deshpande, dr.anirudda.deshpande@gmail.com

Accepted 12 June 2018

DESCRIPTION

A 24-year-old female patient presented with acute-onset, vertex headache associated with photophobia, nausea and vomiting. Initially, the headache was mild (pain scale 4/10), but over the next 4–5 hours, the severity of the headache increased to pain scale of 8/10. She had a history of similar headaches of lesser intensity in the past. There

was also a strong family history of migraine. Her mother and younger sister used to get similar headaches and got relieved with antimigraine medications. Exact details of her medication history were unavailable since the progression of severity of her headache was gradual. This was unlike thunderclap headache and, in high likelihood, another episode of migraine.

Figure 1 MRI brain showing T1-weighted hyperintense pineal gland, as seen on the axial view suggestive of intrapineal gland apoplexy.

Figure 2 Fluid–fluid interface depicted by the horizontal sharp ‘cut-off’ on FLAIR axial view, suggestive of apoplexy, most probably in an intrapineal cyst.

© Author(s) (or their employer(s)). No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Karthik DK, Khardenavis V, Kulkarni S, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-225187

Figure 3 Fluid–fluid interface depicted by the vertical sharp ‘cut-off’ at posterior aspect on T2-weighted sagittal view, suggestive of apoplexy, most probably in an intrapineal cyst.

Figure 4 T1-weighted sagittal view contrast scan suggestive of heterogeneous enhancement of wall of the pineal gland.

Her vitals were stable. Neurological examination, including funduscopy, was grossly normal

On radiological evaluation, the MRI brain showed T1-weighted hyperintense pineal gland, as seen on the axial view suggestive of intrapineal gland apoplexy (figure 1). The fluid–fluid interface is depicted by the horizontal sharp ‘cut-off’ on Fluid Attenuated Inversion Recovery (FLAIR) axial view suggestive of apoplexy, most probably in an intrapineal cyst (figure 2).

Similarly, the vertical sharp ‘cut-off’ on the posterior aspect of the pineal gland on T2-weighted sagittal view (figure 3) was a pointer towards recent haemorrhage probably inside a

pineal cyst. Since the wall of the pineal cyst is thin, it may show heterogeneous enhancement in contrast MRI as shown in T1-weighted sagittal view (figure 4).

Apoplexy (meaning ‘a striking away’ in Greek) is bleeding within internal organs causing associated symptoms. Pineal apoplexy is rare medical condition. It refers to the acute neurological deterioration following bleeding in the pineal region, most commonly into a pineal cyst. It is mostly seen in women, aged between 20 and 30 years. The sensitivity of detection of cysts in the pineal gland depends on the size of the cyst. There is more likelihood of detection of cysts with

Learning points

- ▶ Pineal gland apoplexy is a rare neurological condition and may mimic an episode of migraine headache.
- ▶ MRI in pineal gland apoplexy identifies the intracystic haemorrhage by a characteristic fluid–fluid interface.
- ▶ The differential diagnoses of pineal gland apoplexy are cystic gliomas and pineal parenchymal tumours such as pinealocytomas.

a diameter of more than 5 mm by an MRI scan. Cysts <5 mm in diameter may not be picked up on the MRI.

The differential diagnosis in this case were

- ▶ Cystic gliomas.
- ▶ Pineal parenchymal tumours such as pinealocytomas.¹

The pineal cysts are well-circumscribed, round-shaped lesions anatomically. MR imaging can identify intracystic haemorrhage by a characteristic fluid–fluid interface² However,

pinealocytomas may mimic cysts, and it is difficult to distinguish between the two on routine MRI scan.³

Contributors DKK and AD made substantial contributions to the conception and design of the work, acquisition, analysis and interpretation of the data. VK and SK were involved in drafting the work and revising it critically for important intellectual content. AD approved the final version to be published.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Oosterhout WPJvan, Haan J. Atypical presentation of an apoplexy in a pineal cyst. *J Clin Case Rep* 2015;05:538.
- 2 Sarikaya-Seiwert S1, Turowski B, Hänggi D, Janssen G, Steiger HJ, Stummer W. Symptomatic intracystic hemorrhage in pineal cysts. Report of 3 cases. *J Neurosurg Pediatr* 2009;4:130–6.
- 3 Fakhra S, Escott EJ. Pineocytoma mimicking a pineal cyst on imaging: true diagnostic dilemma or a case of incomplete imaging? *AJNR Am J Neuroradiol* 2008;29:159–63.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow