OPEN ACCESS

CASE REPORT

Aggressive treatment in paediatric or young patients with drug-induced hypersensitivity syndrome (DiHS)/drug reaction with eosinophilia and systemic symptoms (DRESS) is associated with future development of type III polyglandular autoimmune syndrome

Chiyo Morita,^{1,2} Tetsuji Yanase,¹ Tetsuo Shiohara,³ Yumi Aoyama⁴

¹Dermatology, Onomichi General Hospital, Onomichi, Hiroshima, Japan

²Kawasaki Medical School, Kurashiki, Okayama, Japan

³Department of Dermatology, Kyorin University School of Medicine, Tokyo, Japan

⁴Department of Dermatology, Kawasaki Medical School, Kurashiki, Okayama, Japan

Correspondence to

Dr Yumi Aoyama,
ymaoyama@med.kawasaki-m.ac.jp

Accepted 10 August 2018

SUMMARY

We experienced a 6-year-old case of drug-induced hypersensitivity syndrome (DiHS)/drug reaction with eosinophilia and systemic symptoms (DRESS) with subsequent development autoimmune thyroiditis (Hashimoto's thyroiditis), type 1 diabetes with antithyroglobulin, thyroid peroxidase, insulinoma-associated antigen and anti-insulin antibodies at 4 months, alopecia at 7 months, vitiligo, uveitis due to Vogt-Koyanagi-Harada disease at 8 months after clinical resolution of the DiHS/DRESS. He was diagnosed as type III polyglandular autoimmune syndrome (PASIII) after DiHS/DRESS. Prompted by this case, we sought to determine which triggering factors were responsible for later development of PASIII in previously published cases with autoimmune sequelae. In the literature review, five patients with DiHS/DRESS were found to develop autoimmune sequelae consistent with PASIII. All cases with PASIII were much younger than those without them. Four out of the five patients were treated with intravenous immunoglobulin or pulsed prednisolone in the acute stage, although effective in short-term outcomes.

BACKGROUND

Type III polyglandular autoimmune syndrome (PASIII) is a disorder characterised by the coexistence of more than two glandular autoimmune diseases, such as type 1 diabetes.¹ Little information is available on critical triggers in the pathogenesis of PASIII in genetically predisposed individuals. Of note, very frequent component autoimmune diseases of PAS III reportedly occur as sequelae of a distinct type of severe drug eruption, drug-induced hypersensitivity syndrome (DiHS)/drug reaction with eosinophilia and systemic symptoms (DRESS); this syndrome is characterised by sequential reactivations of herpes viruses due to functional defects in regulatory T cells (Tregs).^{2–5} Paediatric DiHS/DRESS is often misdiagnosed as Kawasaki disease due to similar clinical presentations and treated with intravenous immunoglobulin (IVIG).

We describe a child case of DiHS/DRESS with subsequent development of PASIII. In this case, IVIG used would have served to accelerate progression to PASIII, although beneficial in short-term outcomes. Surprisingly, all previously published DiHS/DRESS cases with autoimmune sequelae had clinical features consistent with PASIII and most had been treated with pulsed prednisolone and/or IVIG in the acute stage.

CASE PRESENTATION**Case report**

A 6-year-old boy presented with erythematous lesions on his trunk and extremities. Two months prior to this visit, he suffered a fracture of the right upper arm and received a surgical operation. After the operation, he developed osteomyelitis and was treated with sulfamethoxazole–trimethoprim. Forty-five days later, erythematous plaques began on his extremities and trunk. Sulfamethoxazole–trimethoprim was withdrawn, but his conditions became worse and the patient was referred for further diagnostic workup and treatment to our hospital.

On examination, the patient had numerous 2–5 cm purplish and erythematous plaques on the back and extremities (figure 1A). The patient was hospitalised with fever of 38°C. Physical examination revealed leucocytosis ($14.5 \times 10^9/L$) with atypical lymphocytosis ($2.9 \times 10^9/L$), but no eosinophilia ($0.5 \times 10^9/L$), liver dysfunction (alanine aminotransferase (ALT) 357 U/L) and cervical lymphadenopathy (figure 2). A skin biopsy revealed a spongiotic change of the epidermis with scattered necrotic keratinocytes and a mild perivascular infiltrate. The diagnosis of DiHS/DRESS was made and oral prednisolone, 30 mg/day, was begun. The patient, however, once again experienced the similar erythematous plaques with a high-grade fever and a marked increase in liver enzyme levels (ALT 1857 U/L). On the 12th day of hospitalisation, he received IVIG, at a dose of 1g/kg/day for 2 days given his low immunoglobulin (Ig)G levels (386 mg/dL). After starting IVIG, he showed clearing of

© BMJ Publishing Group Limited 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Morita C, Yanase T, Shiohara T, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-225528

Fig.1

Figure 1 Alterations of clinical presentation of drug-induced hypersensitivity syndrome/drug reaction with eosinophilia and systemic symptoms and autoimmune sequelae. (A) At the initial presentation, the patient had purplish and erythematous plaques on the face. (B) The partial hair loss started 7 months after onset of the disease. (C) Total hair loss associated with generalised vitiligo after 21 months.

lesions and rapid improvement in liver enzyme levels. Because of frequent flaring of liver dysfunction and rashes, we decided to taper prednisolone more gradually over a prolonged period of time (5 mg/day per 2–3 week).

After 4 months, the patient was again admitted to our hospital for nausea, fatigue and excessive thirst. Laboratory tests revealed a marked increase in fasting blood sugar, 580 mg/dL, haemoglobin A1c, 10.2%, anti-insulinoma-associated antigen 2 (IA-2) antibody titre 1.0 U/mL (normal <0.4 U/mL) and anti-insulin antibody titre 50.0 U/mL (<0.4 U/mL). He was diagnosed as having type 1 diabetes mellitus and treatment with insulin was initiated. Because laboratory tests also revealed an increase in antithyroglobulin antibody 36 IU/mL (<28 IU/mL) and antithyroid peroxidase antibody titre 113 IU/mL (<16 IU/mL), he was also diagnosed with Hashimoto's disease. The doses of prednisolone were gradually reduced and eventually withdrawn on day 180.

On days 215–243, he developed alopecia and vitiligo on his back and upper extremities (figure 1B). On day 243, he complained of impaired vision in both eyes. Fundus examination showed multiple areas of neurosensory detachments and focal areas of multiple confluent yellow-coloured placoid choroidal lesions. A diagnosis of concurrent development of Vogt-Koyanagi-Harada disease and prednisolone was again increased to a

dose of 5 mg/day. He developed alopecia totalis with generalised vitiligo 21 months after onset of disease (figure 1C).

INVESTIGATIONS

Prompted by this case, we investigated whether autoimmune diseases with various autoantibodies occurring in reported DiHS/DRESS cases could have clinical features consistent with PASIII. As shown in table 1, five published cases sequentially developing PASIII including this case, were found, although none have reported these autoimmune diseases in the name of PASIII.^{3 5–8} The mean age of these five cases (21 ± 13.4 years) was much younger than that in our 55 institutional cases with DiHS/DRESS (54.5 ± 2.7 years) (manuscript submitted): four of the five cases were categorised as paediatric (<18 years) DiHS/DRESS. Among the three subcategories of PASIII, the present case met the criteria for A and C type of PASIII.¹⁹

DIFFERENTIAL DIAGNOSIS

There are a number of differential diagnostic considerations in children; they include Kawasaki disease and viral eruptions, which show unexplained illness that includes rash, fever and lymphadenopathy. In particular, there is an emerging recognition that some children with DiHS/DRESS have been misdiagnosed

Findings that shed new light on the possible pathogenesis of a disease or an adverse effect

Figure 2 Clinical symptoms and laboratory findings in case 5 with type III polyglandular autoimmune syndrome (present case) after onset of drug-induced hypersensitivity syndrome/drug reaction with eosinophilia and systemic symptoms in relation to treatment. AST, aspartate aminotransferase; ALT, alanine aminotransferase; IVIG, intravenous immunoglobulin; PSL, prednisolone; WCC, white cell count.

as Kawasaki disease and erroneously treated by high doses of IVIG within the first 10 days of fever onset.

TREATMENT

Although clinical experience suggested that high doses of IVIG or pulsed corticosteroids were effective in abrogating the acute inflammation of DiHS/DRESS, our case and previously reported cases with PASIII suggested that these aggressive therapies in paediatric patients paradoxically increased the risk of future development of autoimmune sequelae, particularly PASIII. The potential reward should outweigh the potential risk, especially in a long-term outcome.

OUTCOME AND FOLLOW-UP

Most importantly, pulsed corticosteroids or IVIG was used for treatment of DiHS/DRESS during the acute stage in four of the five reported DiHS/DRESS cases who eventually developed PASIII, except for Case 1, in whom detailed information is not available. Although current literature recommends, DiHS/DRESS cases treated with these aggressive therapies in the acute stage, especially paediatric cases, developed PASIII several months after successful resolution of initial clinical symptoms.⁹⁻¹²

DISCUSSION

Which triggering factors are responsible for later development of PASIII in genetically predisposed individuals after clinical resolution of DiHS/DRESS? Of note, the development of PASIII occurred after administration of a short course of pulsed prednisolone in three of the five reported cases (table 1). Some clues may come from studies of immune reconstitution inflammatory syndrome (IRIS), where a rapid immune recovery on withdrawal or reduction of immunosuppressive agents, paradoxically causes extensive pathology.¹³ Because a rapid and great reduction of prednisolone doses, for example, from 1 g/day to 60 mg/day over

a period of 2 days, is necessarily needed in pulsed prednisolone therapy, thereby increasing the risk of IRIS, pulsed prednisolone therapy for DiHS/DRESS may serve to accelerate aggressive cytotoxic immune responses causing extensive pathology.

Although our patient's young age is unusual in patients with DiHS/DRESS and PASIII, another unusual aspect was its low levels of IgG followed by a marked increase in the levels 1 month later, at which time its levels usually return to normal; this suggests that marked rebound of IgG synthesis may have occurred around the time of IVIG administration in this case: thus, the timing of IVIG administration was likely to coincident with the timing of the rebound and with the peak of liver dysfunction. In this regard, we previously demonstrated that both severe liver damage and defective Treg function in combination could be needed for the subsequent generation of autoantibodies in patients with DiHS/DRESS.¹⁴ In our patient, given the clinical course of the disease, IVIG would have been administered at or immediately after the peak of liver damage and rebound in IgG, A and M synthesis. This timing of IVIG administration may allow autoreactive B cells to function in an uncontrolled fashion: IVIG administration at the peak of severe liver damage could further accelerate the rapid recovery of B cells, thereby causing the subsequent expansions of autoreactive B cells with the specificity to a variety of epitopes. Supporting this hypothesis, De Grandmont *et al* observed that the addition of IVIG in culture of CD40L-stimulated B cells accelerated the differentiation of part of peripheral B cells into IgG-producing plasma cells in the presence of interleukin (IL)-4 and IL-10: importantly, secreted IgGs were shown to be reactive with antigens such as nucleoprotamine, dsDNA, tetanus toxin and human IgG F(ab')₂ fragments. The action of IVIG on DiHS/DRESS may be different depending on the state of activation of the B cells exposed to IVIG.¹⁵

In conclusion, paediatric patients with DiHS/DRESS exhibit an increased risk for subsequently developing PASIII. We

Table 1 Characteristics of patients who developed PAS after drug-induced hypersensitivity syndrome/drug reaction with eosinophilia and systemic symptoms

Patient no. (Ref no.)	Age (y)/ sex	Suspected medications	Systemic corticosteroid therapy (mg/day)	Additional therapy	PAS type	Symptoms of PAS	Autoantibodies	Occurrence of autoimmune sequelae and detection of autoantibodies (weeks or months after onset)
1 ⁶	15/F	Minocycline	PSL 40 mg; escalation dose from 10 mg to 40 mg for 4 days)	None	IIIA	Autoimmune thyroiditis, autoimmune-associated type 1 diabetes	Anti-TPO, TG, TSI, TSHR, GAD, IA2, ANA, Sm and SS-A antibodies	7 weeks: autoimmune thyroiditis in the thyrotoxic phase with anti-TG and TPO antibodies. 5 months: Graves disease with anti-TSI and TSHR antibodies. 7 months: type 1 diabetes with anti-GAD and IA2 antibodies. ANA, anti-Sm and SS-A antibody titres were elevated.
2 ⁷	17/F	Zonisamide	Pulsed PSL; 1 g of mPSL for 3 days followed by 45 mg of PSL	None	IIIA+C	Autoimmune thyroiditis, immune-mediated type 1 diabetes, alopecia, vitiligo. Photophobia, vision impairment, uveitis due to Vogt-Koyanagi-Harada disease	Anti-TPO, TG and TSHR antibodies	2 months: type 1 diabetes, autoimmune thyroiditis (Hashimoto's thyroiditis) with anti-TPO, TG and TSHR antibodies. 5 months: alopecia. 6 months: vision impairment. 9 months: vitiligo. 10 months: Vogt-Koyanagi-Harada disease.
3 ⁸	21/F	Lamotrigine	mPSL 160 mg (40 mg every 6 hours) followed by 80 mg of PSL	IVIg	IIIA	Autoimmune thyroiditis, immune-mediated type 1 diabetes	Anti-TPO, anti-insulin and IA2 antibodies	5 months: type 1 diabetes with anti-insulin and IA2 antibodies. 8 months: autoimmune thyroiditis.
4 ³	46/F	Zonisamide	Pulsed PSL, 1 g of mPSL for 3 days	IVIg	IIIC	Alopecia, autoimmune thyroiditis, Raynaud's phenomenon, sclerodermatous lesions	ANA, RF, anti-ss-DNA, centromere, cardiolipin and TSHR antibodies	6 months: alopecia. 3 years: thyroid dysfunction, Raynaud's phenomenon. 4 years: sclerodermatous lesions with increased titres of ANA.
Five present case	6/M	Sulfamethoxazole/trimethoprim	PSL 36 mg (1.5 mg/kg/day)	IVIg	IIIA+C	Autoimmune thyroiditis, immune-mediated type 1 diabetes, alopecia, vitiligo, uveitis	Anti-TG, TPO, IA-2 and anti-insulin antibodies	4 months: autoimmune thyroiditis (Hashimoto's thyroiditis), type 1 diabetes with anti-TG, TPO, IA-2 and anti-insulin antibodies. 7 months: alopecia. 8 months: vitiligo, uveitis due to Vogt-Koyanagi-Harada disease.

ANA, antinuclear antibodies; F, female; GAD, glutamic acid decarboxylase; IA2, insulinoma-associated antigen-2; IVIg, intravenous immunoglobulin; M, male; PAS, polyglandular autoimmune syndrome; PSL, prednisolone; TG, thyroglobulin; TPO, thyroid peroxidase; TSHR, thyroid stimulating hormone receptor; TSI, thyroid stimulating immunoglobulin.

recommend that patients with DiHS/DRESS, especially paediatric patients, be screened for the generation of various autoantibodies and that at-risk patients be advised to avoid therapies that can accelerate rapid recovery of B cells and T cells, such as IVIG and pulsed prednisolone in the acute stage.

Learning points

- In literature review, five patients with drug-induced hypersensitivity syndrome (DiHS)/drug reaction with eosinophilia and systemic symptoms (DRESS) were found to develop autoimmune sequelae consistent with type III polyglandular autoimmune syndrome (PASIII) 2–7 (average 4.8) months after onset. Four out of the 5 patients were treated with intravenous immunoglobulin or pulsed prednisolone in the acute stage.
- Paediatric or younger patients with DiHS/DRESS have the greatest risk for subsequently developing PASIII, when treated with intravenous immunoglobulin or pulsed prednisolone that can accelerate rapid recovery of B and T cells in the acute stage, although beneficial in the short-term outcome.

Contributors YA and TS contributed to the conception and design, interpretation of data and drafting the article or revising it critically for important intellectual content. All authors contributed to the acquisition of data or analysis. All authors contributed to the final approval of the version published; agreed to be accountable for the article and to ensure that all questions regarding the accuracy or integrity of the article are investigated and resolved.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Kahaly GJ, Frommer L. Polyglandular autoimmune syndromes. *J Endocrinol Invest* 2018;41:91–8.
- Ushigome Y, Kano Y, Ishida T, et al. Short- and long-term outcomes of 34 patients with drug-induced hypersensitivity syndrome in a single institution. *J Am Acad Dermatol* 2013;68:721–8.
- Kano Y, Sakuma K, Shiohara T. Sclerodermoid graft-versus-host disease-like lesions occurring after drug-induced hypersensitivity syndrome. *Br J Dermatol* 2007;156:1061–3.
- Chen YC, Chang CY, Cho YT, et al. Long-term sequelae of drug reaction with eosinophilia and systemic symptoms: a retrospective cohort study from Taiwan. *J Am Acad Dermatol* 2013;68:459–65.
- Aota N, Shiohara T. Viral connection between drug rashes and autoimmune diseases: how autoimmune responses are generated after resolution of drug rashes. *Autoimmun Rev* 2009;8:488–94.
- Brown RJ, Rother KI, Artman H, et al. Minocycline-induced drug hypersensitivity syndrome followed by multiple autoimmune sequelae. *Arch Dermatol* 2009;145:63–6.
- Hirakawa A, Kobayashi H, Takasaka M, et al. A case of drug induced hypersensitivity syndrome triggering various autoimmune diseases. *Clin Dermatol* 2016;70:391–6.
- Singer EM, Wanat KA, Rosenbach MA. A case of recalcitrant DRESS syndrome with multiple autoimmune sequelae treated with intravenous immunoglobulins. *JAMA Dermatol* 2013;149:494–5.
- Ben-Skowronek I, Michalczyk A, Piekarski R, et al. Type III polyglandular autoimmune syndromes in children with type 1 diabetes mellitus. *Ann Agric Environ Med* 2013;20:140–6.
- Teng P, Tan B. Carbamazepine-induced DRESS syndrome in a child: rapid response to pulsed corticosteroids. *Dermatol Online J* 2013;19:18170.
- Comfere NI, Sartori-Valinotti JC, Bruce AJ, et al. Successful treatment of lamotrigine-associated drug hypersensitivity syndrome with intravenous IgG. *J Am Acad Dermatol* 2012;66:e249–e250.
- Marcus N, Smuel K, Almog M, et al. Successful Intravenous Immunoglobulin Treatment in Pediatric Severe DRESS Syndrome. *J Allergy Clin Immunol Pract* 2018;6.
- Shiohara T, Kurata M, Mizukawa Y, et al. Recognition of immune reconstitution syndrome necessary for better management of patients with severe drug eruptions and those under immunosuppressive therapy. *Allergol Int* 2010;59:333–43.
- Takehara A, Aoyama Y, Kurosawa M, et al. Longitudinal analysis of antibody profiles against plakins in severe drug eruptions: emphasis on correlation with tissue damage in drug-induced hypersensitivity syndrome and drug reaction with eosinophilia and systemic symptoms. *Br J Dermatol* 2016;175:944–52.
- de Grandmont MJ, Racine C, Roy A, et al. Intravenous immunoglobulins induce the in vitro differentiation of human B lymphocytes and the secretion of IgG. *Blood* 2003;101:3065–73.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow