OPEN ACCESS

CASE REPORT

Invasive mediastinal aspergillosis presenting as superior vena cava syndrome in an immunocompetent patient

Yasmin Rahim,¹ Aisha Memon,² Javaid Ahmed Khan¹

¹Department of Pulmonology and Critical Care, Aga Khan University Hospital, Karachi, Pakistan

²Department of Histopathology, Aga Khan University Hospital, Karachi, Pakistan

Correspondence to
Dr Yasmin Rahim,
yrahim80@hotmail.com

Accepted 15 August 2018

SUMMARY

Invasive aspergillosis (IA) is a disease of the immunocompromised with a predilection for the lungs, although dissemination to all organs is possible. Its diagnosis remains a challenge due to the absence of specific clinical manifestations and laboratory findings. In most cases, diagnosis is eventually made via invasive methods. It carries with it a high mortality due to late diagnosis and delayed treatment. Here, we report a fascinating case of a young, otherwise healthy, immunocompetent patient that presented to us with superior vena cava syndrome and a mediastinal mass. It was anticipated that a malignancy would be found on further workup but, in fact, what was eventually discovered was a case of IA. Our report accentuates the significance of including IA as a differential while diagnosing a mediastinal mass in an immunocompetent host as patient outcome is determined by timely diagnosis and treatment.

BACKGROUND

Aspergillosis comprises a spectrum of diseases related to host factors.¹⁻³ Invasive aspergillosis (IA) is a disease of concern due to high mortality in transplant recipients and patients with haematological malignancies. Although rare, its occurrence has been reported in immunocompetent hosts, with multisystem involvement.⁴⁻⁶ Here, we present a

Figure 2 Chest X-ray showing widened mediastinum, elevated right hemidiaphragm and blunting of the right costophrenic angle.

case of a young immunocompetent man presenting with signs of superior vena cava (SVC) obstruction and a mediastinal mass. Suspicion of a malignancy was high but on biopsy was found to have IA of the mediastinum.

CASE PRESENTATION

A 28-year-old man was referred to our pulmonary outpatient clinic (Aga Khan University Hospital, Karachi) for the workup of a mediastinal mass. He had a 12-month history of right-sided chest pain,

Figure 1 Right anterior neck and chest wall showing prominent neck and chest wall veins and mediastinotomy scar.

Figure 3 CT of the chest. (A) Lung window showing right anterior mediastinal mass with right-sided effusion. (B) Mediastinal window showing mediastinal mass occluding right main bronchus. (C) Coronal section showing invasion of the mediastinal structures.

© BMJ Publishing Group Limited 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Rahim Y, Memon A, Khan JA. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-225614

Figure 4 Periodic acid-Schiff stain highlighting mostly fungal spores, rare hyphae seen.

with radiation to the back. He also complained of shortness of breath, reduced appetite, weight loss, difficulty swallowing and swelling of his face and neck for the past 5 months. He denied fever, night sweats, cough or past tuberculosis (TB) contact. He had a history of smoking in the distant past, that is, 2 years of smoking around 10 years ago. He denied alcohol and illicit drug use. He sought medical attention 6 months after his symptoms began in his hometown. He was started on treatment for asthma, that is, inhaled corticosteroids and inhaled long-acting beta-agonist and a 3-week tapering dose of oral prednisolone (starting at 10 mg twice a day) with no improvement whatsoever. A chest X-ray (CXR) was ordered and a suspicious finding on that led to a CT of the thorax, which showed a mediastinal mass. Despite two negative sputum samples for acid-fast bacilli (AFB), he was

Figure 5 Periodic acid-Schiff stain highlighting mostly fungal spores, rare hyphae seen.

Figure 6 Gomori methenamine-silver stain, highlighting fungal bodies.

started on empiric antituberculous therapy. This was discontinued after 3 weeks as there was no clinical improvement.

He was then referred to us for further workup and management with a strong suspicion of malignancy (likely lymphoma).

On initial assessment, vital signs were within normal values. There was evidence of congested neck, prominent chest wall veins and facial plethora (figure 1).

Chest examination revealed absent breath sounds on the lower one-third of the right posterior chest and an otherwise unremarkable cardiovascular and systemic examination.

INVESTIGATIONS

Laboratory tests were normal. CXR was concerning for a widened mediastinum and right-sided effusion (figure 2).

Diagnostic thoracentesis revealed a lymphocyte dominant exudate, a white cell count of $6.673 \times 10^9/L$ (lymphocytes 95%), protein of 4.8 g/dL, glucose of 88 mg/dL and Lactate dehydrogenase (LDH) of 323 U/L. Pleural fluid cytology revealed no malignant cells. Serological testing for HIV was negative. Serum immunoglobulin (IgG, IgA, IgM) levels were within normal range.

CT thorax showed an $11 \times 9 \times 8$ cm mass centred in the anterior and middle mediastinum, more on the right side. It showed invasion of the right atrium, right pulmonary artery with partial

Figure 7 *Aspergillus* colonies.

invasion of the left atrium and left inferior pulmonary vein. Partial obliteration of the right main stem bronchus was noted along with narrowing of the SVC and prominent collaterals (figure 3A–C).

OUTCOME AND FOLLOW-UP

He underwent anterior mediastinotomy, with mediastinal mass biopsy, revealing non-necrotising granulomas, with multinucleated giant cells and histiocytes. Scant fragments of fibrocollagenous tissue noted. Periodic acid-Schiff and Gomori methenamine-silver stains highlighted occasional septate fungal hyphae (figures 4, 5 and 6). Mycology cultures grew the *Aspergillus flavus* complex (figure 7). Mediastinal tissue AFB smear and culture were negative.

Therapy with oral voriconazole 400 mg twice a day for 1 day followed by 200 mg twice a day was started after histological confirmation, and the patient was asked to follow up in 4 and then 8 weeks, while continuing therapy. Patient showed marked improvements in symptoms with progressive weight gain and increased oral intake. Repeat CT scan is planned at 6 months to look for mass resolution.

DISCUSSION

To the best of our knowledge, this is the second case of SVC obstruction secondary to IA of the mediastinum in an immunocompetent host published in the literature. Association with other infectious diseases namely syphilis, TB, histoplasmosis and actinomycosis have been described.

Aspergillus is a spore-forming saprophytic fungus, ubiquitous to the environment. With approximately 200 species, *Aspergillus fumigatus* is the most commonly isolated pathogen.

Aspergillus species cause a broad spectrum of disease, ranging from hypersensitivity reactions to direct angioinvasion. The lung is the most common site of infection with disease ranging from allergic bronchopulmonary aspergillosis, chronic necrotising pneumonia, aspergilloma and IA.⁶

IA is primarily a disease of immunocompromised patients, with prolonged neutropenia constituting a predominant risk factor,⁷ followed by haematopoietic stem cell transplants or solid organ transplants (especially lung transplantation) and advanced AIDS.^{8,9} In the above population of patients, host factors govern the risk and clinicopathological features of invasive disease.^{1,2} Rarely invasive pulmonary aspergillosis may complicate chronic obstructive pulmonary disease, particularly when treated with oral corticosteroids (OCS), bacterial or viral pneumonia, sepsis, liver failure, alcoholism or sinusitis.^{10–12}

Our patient's laboratory findings revealed no evidence of neutropenia or immunosuppression, raising a suspicion of whether his 2-year smoking history or 3-week history of OCS use could have led to some abnormalities in the pulmonary defence mechanism. This seemed an unlikely possibility as the patient was symptomatic months before inhaled or OCS therapy was prescribed.

The respiratory tract remains the primary focus of disease, with poor disease control leading to mediastinal extension and haematogenous dissemination that can involve any organ.¹³

The diagnosis of IA remains a challenge due to the absence of specific clinical manifestations, non-pathognomonic radiological findings and poor sensitivity of sputum cultures. In most cases, diagnosis is made by tissue isolation through invasive methods, as was the case in our patient. However, we felt attempting a less invasive approach namely image-guided fine-needle aspiration would have yielded the same results and

should have been attempted first. Serological tests are adjuncts to support or exclude the diagnosis in the appropriate clinical context; however, they cannot be solely relied on for a definitive diagnosis.

Published literature highlights the poor, usually fatal outcome in patients with disease involving the mediastinum, despite initiation of antifungal therapy.^{14–16} This particularly holds truth in cases with delayed diagnosis leading to treatment delay and fatal outcomes. In this case, the patient continues to have a favourable outcome, with ongoing follow-up.

Treatment is often prolonged, with the duration of treatment determined empirically, depending on the extent of IA, the response to therapy and the immune status of the patient. Voriconazole is the first-line therapy for IA, with treatment duration depending on disease burden and host immunocompetence. The Infectious Diseases Society of America recommends 6 to 12 weeks of therapy for pulmonary aspergillosis.¹⁶ In immunosuppressed patients, therapy should be continued throughout the period of immunosuppression and until the resolution of lesions.¹⁶ Surgical treatment is reserved for patients who do not tolerate antifungal therapy or have residual localised but active lesions, despite adequate antifungal therapy.

This case highlights the importance of including the suspicion of IA in the differential diagnosis of a mediastinal mass presenting with signs of SVC syndrome irrespective of immune status, where TB and malignancy represent valid possibilities, as disease outcome is directly linked with early diagnosis and therapy.¹⁷ Early and timely treatment prevents future complications and death.

Learning points

- Invasive aspergillosis (IA) is a rare but important consideration in immunocompetent patients presenting with signs of superior vena cava obstruction and a mediastinal mass.
- Suspicion of IA in a patient, who does not respond to antibiotic or antituberculous therapy, is imperative in early diagnosis, allowing timely initiation of aggressive antifungal therapy and increasing the chances of a favourable outcome.
- Imaging-guided percutaneous needle biopsy of mediastinal masses should be attempted first before reverting to a more invasive approach namely surgery.

Contributors The case report was conceptualised by JAK. YR reviewed patient records and examination along with literature research and writing the case report. Histopathology images were provided by AM. The final case report was approved by JAK.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Chamilos G, Luna M, Lewis RE, *et al*. Invasive fungal infections in patients with hematologic malignancies in a tertiary care cancer center: an autopsy study over a 15-year period (1989–2003). *Haematologica* 2006;91:986–9.

- 2 Shaukat A, Bakri F, Young P, *et al*. Invasive filamentous fungal infections in allogeneic hematopoietic stem cell transplant recipients after recovery from neutropenia: clinical, radiologic, and pathologic characteristics. *Mycopathologia* 2005;159:181–8.
- 3 Siddiqui S, Anderson VL, Hilligoss DM, *et al*. Fulminant mulch pneumonitis: an emergency presentation of chronic granulomatous disease. *Clin Infect Dis* 2007;45:673–81.
- 4 Allam MF, Del Castillo AS, Díaz-Molina C, *et al*. Invasive pulmonary aspergillosis: identification of risk factors. *Scand J Infect Dis* 2002;34:819–22.
- 5 Han K-H, Kim J-H, Shin SY, *et al*. A case of invasive pulmonary aspergillosis with direct invasion of the mediastinum and the left atrium in an immunocompetent patient. *Tuberc Respir Dis* 2014;77:28–33.
- 6 Shakoor MT, Ayub S, Ayub Z, *et al*. Fulminant invasive aspergillosis of the mediastinum in an immunocompetent host: a case report. *J Med Case Rep* 2012;6:311.
- 7 Gerson SL, Talbot GH, Hurwitz S, *et al*. Prolonged granulocytopenia: the major risk factor for invasive pulmonary aspergillosis in patients with acute leukemia. *Ann Intern Med* 1984;100:345–51.
- 8 Kotloff RM, Ahya VN, Crawford SW. Pulmonary complications of solid organ and hematopoietic stem cell transplantation. *Am J Respir Crit Care Med* 2004;170:22–48.
- 9 Soubani AO, Miller KB, Hassoun PM. Pulmonary complications of bone marrow transplantation. *Chest* 1996;109:1066–77.
- 10 Segal BH. Aspergillosis. *N Engl J Med* 2009;360:1870–84.
- 11 Stevens DA, Melikian GL. Aspergillosis in the 'nonimmunocompromised' host. *Immunol Invest* 2011;40(7-8):751–66.
- 12 Zmeili OS, Soubani AO. Pulmonary aspergillosis: a clinical update. *QJM* 2007;100:317–34.
- 13 Hendrix WC, Arruda LK, Platts-Mills TA, *et al*. Aspergillus epidural abscess and cord compression in a patient with aspergilloma and empyema. *Am Rev Respir Dis* 1992;145:1483–6.
- 14 Parameswaran K, Joshi M, Ravindran P. Unusual radiological presentation and rapid fatal progression of invasive pulmonary aspergillosis in an immunocompetent young patient. *Respirology* 1999;4:287–90.
- 15 Taouli B, Cadi M, Leblond V, *et al*. Invasive aspergillosis of the mediastinum and left hilum: CT features. *AJR Am J Roentgenol* 2004;183:1224–6.
- 16 Walsh TJ, Anaissie EJ, Denning DW, *et al*. Treatment of aspergillosis: clinical practice guidelines of the Infectious diseases society of America. *Clin Infect Dis* 2008;46:327–60.
- 17 Cook DJ, Achong MR, King DE. Disseminated aspergillosis in an apparently healthy patient. *Am J Med* 1990;88:74–6.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow