Laws of attraction: management of magnetic foreign body ingestion

Michael Nicoara, Shinban Liu, George Ferzli

General Surgery, NYU Langone Medical Center, Brooklyn, New York, USA

Correspondence to
Dr Shinban Liu,
shinban.liu@nyumc.org

MN and SL contributed equally.

Accepted 21 June 2018

DESCRIPTION

An 11-year-old girl with global developmental delay presents with a 4-day history of nausea, vomiting and abdominal pain after the ingestion of multiple foreign bodies. She was haemodynamically stable with no gross abnormalities on laboratory tests. An abdominal X-ray was obtained which demonstrated two radio-opaque foreign bodies in the left lower quadrant without free air (*figure 1A*). On physical examination the patient had significant point tenderness in the left lower quadrant with guarding. The decision was made to proceed to the operating room where a diagnostic laparoscopy was performed. On entering the abdomen, there was no significant free fluid or gross contamination. After running the bowel, two magnets were identified in adjacent loops of small intestine with evidence of bowel wall ischaemia in-between. Approximately 15 cm of jejunum was inspected and the foreign bodies which included two magnets and an earring (*figure 1B*) were safely delivered with primary anastomosis. The patient was discharged home on post-operative day 2 after regaining bowel function and tolerating diet.

In the USA, over 100 000 cases of foreign body ingestion are reported yearly, with 80% of them

occurring in children.¹ Most foreign bodies pass spontaneously. However, about 15% of cases will require endoscopic intervention and <1% require surgical intervention. Commonly ingested foreign bodies include coins, toys, bones, pins and magnets. Toys and other appliances containing high-powered magnets made of neodymium are becoming more common in the household. Patients who have ingested these items are often asymptomatic but may present with pain or an acute abdomen if perforation has ensued.

Management of foreign bodies in a symptomatic patient requires prompt exploration and removal. This can be performed endoscopically or surgically depending on the level of involvement and extent of contamination. If the objects are present in the oesophagus, stomach or proximal small bowel, endoscopy should be performed to retrieve the objects and examine the organ structures for injury. In asymptomatic patients with objects beyond the proximal small bowel, expectant management may be performed with serial imaging to evaluate the transit of the foreign body.² The ingestion of multiple magnets presents a unique situation in which foreign objects are not able to pass with normal motility due to tethering between loops of bowel, which in turn

Figure 1 (A) Abdominal X-ray demonstrating multiple foreign objects in the small bowel without free air. (B) Gross specimen with two magnetic foreign bodies and an earring removed from the small bowel.

© BMJ Publishing Group Limited. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Nicoara M, Liu S, Ferzli G. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-225939

may become ischaemic and perforate.³ Surgical exploration may be performed laparoscopically or by an open approach depending on the situation and operator's experience.

Learning points

- ▶ In an asymptomatic patient, foreign object ingestion may be treated with expectant management and serial imaging.
- ▶ Magnetic foreign bodies may cause tissue ischaemia and perforation if adjacent loops of bowel are involved.
- ▶ Intervention can be pursued endoscopically or surgically depending on the level of involvement and extent of contamination.

Contributors MN, SL and GF contributed equally to the diagnosis and treatment of the patient, acquisition of data, interpretation and reporting of the case.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Wyllie R. Foreign bodies in the gastrointestinal tract. *Curr Opin Pediatr* 2006;18:563–4.
- 2 Liu S, de Blacam C, Lim FY, *et al*. Magnetic foreign body ingestions leading to duodenocolonic fistula. *J Pediatr Gastroenterol Nutr* 2005;41:670–2.
- 3 Hussain SZ, Bousvaros A, Gilger M, *et al*. Management of ingested magnets in children. *J Pediatr Gastroenterol Nutr* 2012;55:239–42.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow