



OPEN ACCESS

Advanced Lung Disease and Lung Transplantation, Inova Fairfax Hospital, Falls Church, Virginia, USA

Correspondence to
Dr Christopher King,
christopher.king@inova.org

Accepted 22 September 2018

Fungal thyroiditis in a lung transplant recipient

Christopher King, Stephen Clement, Shalika Katugaha, Anne Whitney Brown

DESCRIPTION

A 59-year-old man was admitted with 3 weeks of worsening shortness of breath 18 months after receiving




Figure 1 Chest CT showing multiple pulmonary nodules in the right lower lung field (red arrows).




Figure 2 Fungal hyphae are present on Gomori Methenamine-Silver (GMS) stain from CT-guided lung biopsy (x200).



Figure 3 CT of the neck revealed a mass-like lesion in the right lobe of the thyroid (green arrow).

a bilateral lung transplant for idiopathic pulmonary fibrosis. His immunosuppression included tacrolimus, everolimus and low-dose prednisone with no antifungal prophylaxis at the time of admission. CT chest revealed multiple, bilateral pulmonary nodules (figure 1—red arrows). CT-guided biopsy revealed fungal hyphae (figure 2). The initial CT and ultrasound of the neck at the onset of sore throat was negative; however, repeat CT neck for evolving neck pain and dysphasia during hospital course showed a mass-like lesion in the right thyroid lobe with extensive surrounding inflammatory changes (figure 3—green arrow). The lesion was also visualised on ultrasound, where it appeared as a hypoechoic solitary nodule (figure 4). Laboratory evaluation revealed hyperthyroidism, with a Thyroid Stimulating Hormone (TSH) of $<0.01 \mu\text{U/mL}$ (normal: $0.35\text{--}4.94 \mu\text{U/mL}$) and a free T4 of 2.72 ng/dL (normal: $0.70\text{--}1.48 \text{ ng/dL}$). The patient was initiated on dual antifungal therapy with liposomal amphotericin and posaconazole, resulting in rapid resolution of neck pain. However, repeat CT neck 2 weeks after revealed a thyroid abscess in the




Figure 4 Thyroid ultrasound reveals a $2.8 \times 2.3 \text{ cm}$ hypoechoic lesion in the right thyroid isthmus.




Figure 5 Fungal hyphae were present in the necrotic material from the thyroid abscess debridement (H&E, x400).



© BMJ Publishing Group Limited 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: King C, Clement S, Katugaha S, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-227033

Images in...

right lobe, and subsequently the patient underwent right thyroid lobectomy and isthmusectomy. Surgical debridement of the thyroid abscess revealed fungal hyphae; thus, confirming the diagnosis of fungal thyroiditis (figure 5). A specific fungal pathogen was never successfully cultured from our patient. He was treated for presumed

Aspergillus infection, the most common etiology of fungal thyroiditis, with clinical improvement.¹

Acknowledgements Haresh Mani, MD Min Kim, MD.

Contributors CK, SK and AWB: planning, writing, revision. SC: planning, revision.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCE

- 1 Goldani LZ, Zavascki AP, Maia AL. Fungal thyroiditis: an overview. *Mycopathologia* 2006;161:129–39.

Learning points

- ▶ Thyroiditis is a rare manifestation of disseminated fungal infection in immunocompromised hosts.
- ▶ *Aspergillus* spp are the most commonly implicated pathogen, although there are reports of cases due to *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Coccidioides immitis* and *Candida* spp.
- ▶ Antifungal therapy targeting the culprit organism and consideration of surgical debridement are the mainstays of treatment.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow