Perineurioma of the sigmoid colon

Guled M Jama,¹ Matthew Evans,² Muhammad W Fazal,³ Deepak Singh-Ranger¹

¹Department of General Surgery, The Royal Wolverhampton NHS Trust, Wolverhampton, West Midlands, UK

²Department of Histopathology, The Royal Wolverhampton NHS Trust, Wolverhampton, West Midlands, UK

³Department of Gastroenterology, The Royal Wolverhampton NHS Trust, Wolverhampton, West Midlands, UK

Correspondence to

Dr Guled M Jama, guled.jama@nhs.net

Accepted 12 September 2018

DESCRIPTION

A 24-year-old man presented to the surgical outpatient clinic with a 1-year history of intermittent, painless, fresh rectal bleeding associated with defaecation. The blood was noticeable as streaks on the toilet paper. His bowel habit remained otherwise unchanged. He reported no weight loss or family history of bowel cancer. His medical history was unremarkable.

On examination, there were no clinical features of anaemia, jaundice, cyanosis, clubbing, oedema or cervical lymphadenopathy. Proctoscopy revealed an internal haemorrhoid at the 7 o'clock position. Blood test results included a haemoglobin level of 158 g/L, mean corpuscular volume of 87.0 fL and mean corpuscular haemoglobin concentration of 331 g/L, all of which were within normal limits.

On flexible sigmoidoscopy, a 3 mm polyp within the sigmoid colon was identified. This was a well-circumscribed, flat elevated lesion (Paris classification type 0-IIa) (figure 1). Its surface features on narrow band imaging were not typical of an adenomatous or hyperplastic polyp. Given its unusual appearance, the polyp was removed.

Histology showed it to be a bland mesenchymal lesion of the colonic wall. The lamina propria appeared to be expanded by a vaguely nodular proliferation of spindle cells percolating between distorted non-serrated crypts (figure 2A). There was no cytological atypia, mitotic activity or necrosis. Very small blood vessels were scattered throughout. There was no perivascular condensation or eosinophilia to suggest an inflammatory fibroid polyp. Immunohistochemical staining showed widespread although weak positivity for epithelial membrane antigen (figure 2B). DOG1, CD34, CD117, smooth muscle actin, desmin, S100 and cytokeratin AE1/3 were all negative. The features were consistent with a perineurioma.

Colonic perineuriomas, or fibroblastic polyps, are rare benign mucosal lesions comprising a proliferation of bland spindle cells expressing markers of

Figure 2 Photomicrograph. (A) Nodular spindle cell lesion percolating between distorted but non-serrated colonic crypts (H&E stain, $\times 100$ magnification). (B) Weak staining in the spindle cells (epithelial membrane antigen stain, $\times 200$ magnification).

perineurial differentiation.¹ They are usually found in the rectum and sigmoid colon, although they have been reported proximal to the splenic flexure.² With an estimated incidence of 0.1%–1.46% of all colonic polyps, they tend to be discovered incidentally in patients undergoing routine screening colonoscopy.

Endoscopically, they appear as solitary, sessile, well-circumscribed mucosal lesions and range in size from 0.2 cm to 1.5 cm. Characteristic histological features include spindle cell proliferation within the lamina propria which often separates and distorts the colonic crypt architecture.

Two variants are recognised: those with and those without accompanying serrated epithelium; the former may superficially resemble hyperplastic polyps or sessile serrated lesions. The pathogenesis of the perineurial-type stroma is poorly understood. There is increasing evidence to suggest that the epithelial component of serrated fibroblastic polyps harbours BRAF mutations which may trigger epithelial–mesenchymal interactions, causing fibroblast transdifferentiation and proliferation. The development of the less common non-serrated variant, of which this case is an example, has yet to be determined.³ It is possible that the two variants represent entirely separate biological entities.

Figure 1 Flexible sigmoidoscopy. Gross appearance of the polyp under white light.

Learning points

- ▶ Colonic perineuriomas are rare benign mucosal lesions; however, their characterisation is important in determining colonoscopic surveillance.
- ▶ Decisions about follow-up surveillance colonoscopy should be based on the nature and size of the accompanying serrated epithelial component (if present), according to existing guidelines for serrated colorectal polyps in general.
- ▶ There are no reported cases of recurrence or metastases.

© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Jama GM, Evans M, Fazal MW, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-227170

To our knowledge, there have been no reported cases of recurrence or metastases of colonic perineuriomas. However, the neoplastic nature of the serrated variant means that they should be classified based on the criteria used for serrated colorectal polyps in general, making postpolypectomy surveillance in some cases appropriate.³

Contributors GMJ performed the literature review and wrote the manuscript. ME and MWF provided the images and wrote the manuscript. DS-R reviewed and edited the manuscript. All authors approved the final version of the manuscript before submission.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Hornick JL, Fletcher CD. Intestinal perineuriomas: clinicopathologic definition of a new anatomic subset in a series of 10 cases. *Am J Surg Pathol* 2005;29:859–65.
- 2 van Wyk AC, van Zyl H, Rigby J, et al. Colonic perineurioma (benign fibroblastic polyp): case report and review of the literature. *Diagn Pathol* 2018;13:16.
- 3 Erlenbach-Wünsch K, Bihl M, Hartmann A, et al. Serrated epithelial colorectal polyps (hyperplastic polyps, sessile serrated adenomas) with perineurial stroma: clinicopathological and molecular analysis of a new series. *Ann Diagn Pathol* 2018;35:48–52.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow