Giant craniopharyngioma in an adult presenting with new onset seizure

Subhanudh Thavaraputta,¹ Ana Marcella Rivas²

¹Department of Internal Medicine, Texas Tech University Health Sciences Center, Lubbock, Texas, USA

²Division of Endocrinology, Department of Internal Medicine, Texas Tech University Health Sciences Center, Lubbock, Texas, USA

Correspondence to

Dr Ana Marcella Rivas, marcella.rivas@ttuhsc.edu

Accepted 4 September 2018

DESCRIPTION

A 43-year-old man presented to the emergency department after experiencing a first episode seizure. No further information was available at presentation as the patient was confused on his arrival. On physical examination, vital signs were normal. The patient was confused, and his Glasgow Coma Score was 12(E3V4M5). Initial workup included complete blood count, electrolytes and ECG were normal. Thyroid function tests, insulin growth factor-1 and plasma random cortisol level were within the normal range. Testosterone levels were low, and luteinising hormone was inappropriately normal. CT demonstrated a cystic mass with peripheral calcification measuring 6.0×4.5×3.9 cm. The lesion originated from the pituitary fossa and expanded superiorly distorting the third ventricle and the left lateral ventricle causing hydrocephalus (figure 1). A pituitary MRI confirmed these findings (figure 2). The patient underwent

Figure 2 Sagittal (A) and coronal (B) sections of T1 MRI of the head demonstrates a cystic mass with minimal enhancing solid components and calcifications (white arrow) in the sella region with suprasellar extension. The mass measures 6.01×4.52×4.07 cm and extends superiorly into the third ventricle, the body of the left lateral ventricle and right parasellar causing hydrocephalus.

Figure 1 Sagittal (A) and axial (B) sections of CT scan of the head demonstrate a cystic mass lesion with peripheral calcification (black arrows), measuring 6.0×4.5×3.9 cm, originating from the pituitary gland, expanding into the suprasellar cistern, interpeduncular cistern and distorting the floor of the third ventricle with significant compromise of foramen of Monroe, resulting in bilateral ventricular dilatation.

a craniotomy with mass resection. The procedure was uneventful. Pathology demonstrated nodular whorls and irregular trabeculae of well-differentiated squamous epithelium, bordered by palisading columnar cell consistent with an adamantinomatous craniopharyngioma (CP) (WHO grade I) (figure 3). Postoperatively, the patient developed panhypopituitarism, and on close monitoring, he was noted to have hypothalamic dysfunction manifesting with hypothermia, hypoglycaemia and adipsia.

CPs are benign, slow-growing epithelial tumours that derive from ectodermal remnants of Rathke's pouch. They represent 2%–5% of all primary intracranial tumours with an incidence rate of 1–2 cases per million a year. About 30%–40% of cases present during adolescence, while around 25% are diagnosed in patients over the age of 25.¹ Over 60% of cases arise from within the sellar and presenting symptoms result from compression to the surrounding structures. Clinical manifestations include visual defects, pituitary insufficiency, ocular motor nerve palsies and symptoms of increased intracranial pressure. Two pathological types of CP are recognised: adamantinomatous and papillary CP. Adamantinomatous CPs can be seen in patients of all ages while papillary CP are more common in adults.¹ On imaging, calcification of a tumour is seen in 60%–80% of CP. The mean average tumour size at the time of presentation is 3 cm in adults² and 5 cm in children.³

Giant CPs, described as a tumour larger than 5–6 cm on their largest axis, are more common in children. They are usually found as adamantinomatous CP and can range from 5 to 11 cm.² In adults,

© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Thavaraputta S, Rivas AM. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-227244

Figure 3 H&E stain of pituitary mass demonstrated (A) nodular whorls, lobules and irregular of well-differentiate trabecular squamous epithelium bordered by palisaded columnar epithelium. (B) There are multiple nodules of wet keratin (black asterisk) and stellate reticulum (black arrow) which are classic histological feature of the adenomatous craniopharyngioma.

giant CPs are extremely rare and infrequently reported in the literature. Large tumours are associated with worse prognosis due to severe mass effect and residual neurological sequelae. Seizures as a manifestation of pituitary lesions are rare, but more commonly seen with large tumours. The mechanism triggering seizures is not fully understood. It has been suggested that it may relate to mass effect exerted over the suprasellar and/or temporal areas, eliciting an epileptic discharge. Regardless of the tumour size, CPs rarely undergo malignant transformation.¹ Treatment includes resection and possible adjuvant chemotherapy.

Learning points

- ▶ Craniopharyngiomas are usually benign and their morbidity usually results from damage of surrounding structures due to compression.
- ▶ In adults, the average tumour size is 3 cm, and giant craniopharyngiomas are rare.
- ▶ Larger tumours are more likely to compress surrounding structures and associated with worse outcomes.

Contributors ST: first author, patient care, data acquisition, drafting manuscript, manuscript revision, gathering information and literature search. AMRM: patient care, manuscript revision, data acquisition, final approval and corresponding author.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Melmed S, Kleinberg D. Chapter 9 - Pituitary masses and tumors. Melmed S, *Williams textbook of endocrinology*. 13 edn: Philadelphia, 2016:232–99.
- 2 Al-Mefty O, Hassounah M, Weaver P, *et al*. Microsurgery for giant craniopharyngiomas in children. *Neurosurgery* 1985;17:585–95.
- 3 Elliott RE, Jane JA, Wisoff JH. Surgical management of craniopharyngiomas in children: meta-analysis and comparison of transcranial and transsphenoidal approaches. *Neurosurgery* 2011;69:630–43.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow