Red colour venous flow in the suprasternal view: a red flag sign

Sahil Mahajan,¹ Gopal Chandra Ghosh,¹ Oommen K George²

¹Department of Cardiology, Christian Medical College and Hospital Vellore, Vellore, Tamil Nadu, India

²Department of Cardiology, Christian Medical College and Hospital Vellore, Vellore, Tamil Nadu, India

Correspondence to
Dr Gopal Chandra Ghosh,
gcghosh86@gmail.com

Accepted 18 September 2018

DESCRIPTION

A 3-day-old full-term neonate, born by normal vaginal delivery, was presented with respiratory distress and mild cyanosis. His room air saturation in all four limbs was 84%. The clinical examination was unremarkable. Chest X-ray showed cardiomegaly and increased pulmonary vascular markings. Transthoracic echocardiography revealed 8 mm ostium secundum atrial septal defect (ASD) with bidirectional shunt and dilated right atrium and right ventricle with absence of individual pulmonary veins (PV) draining into left atrium (LA). Suprasternal view showed vertical vein (VV) which was recognised as a vessel lateral to LA with red-coloured continuous flow on Doppler (figure 1). The PVs were draining into the venous confluence behind the LA which was draining into the VV. VV connects into the left innominate vein which finally forms the right superior vena cava (figure 1, video 1).

Clinical presentation of total anomalous pulmonary venous connection (TAPVC) varies from congestive heart failure at one end of spectrum to deep cyanosis at the other which in turn depends on the balance between the interatrial communication and pulmonary venous outflow obstruction.¹ Our case highlights the importance of recognising additional channel in relation to aorta in suprasternal view. Usually there is only one vessel visible except aortic arch, which is right pulmonary artery. The differential diagnosis of additional vessels can be aortopulmonary collaterals, patent ductus arteriosus (PDA) arising from aortic arch, VV in TAPVC and retroaortic innominate vein.² Meticulous echocardiography can be rewarding in differentiation of these structures. Aortopulmonary collaterals and PDA will show high-velocity continuous flow.

Video 1 Video demonstrating pulmonary venous drainage into a common chamber which finally drains into a vertical vein.

Retroaortic innominate vein will show low-velocity venous flow (blue flow) towards the heart while VV can be identified by low-velocity venous flow away from the heart (red flow). Our patient underwent successful rechanneling of PVs to the LA with ligation of VV and closure of ASD.

Learning points

- ▶ Supracardiac variety of total anomalous pulmonary venous connection (TAPVC) is the most common type of TAPVC.
- ▶ Transthoracic echocardiographic suprasternal view is the ideal view for anatomical delineation of pulmonary venous drainage.
- ▶ Demonstration of vertical vein is the cornerstone for diagnosis and meticulous echocardiography helps in differentiating it from other vascular structure.

Contributors SM drafted the manuscript. GCG and OKG supervised in writing.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Lee ML, Wu MH, Wang JK, *et al.* Echocardiographic assessment of total anomalous pulmonary venous connections in pediatric patients. *J Formos Med Assoc* 2001;100:729–35.
- 2 Sukulal K, Bijulal S, Tharakan JA. Differential diagnosis of vascular structures in relation to upper ascending aorta: The retro-aortic innominate vein. *Ann Pediatr Cardiol* 2013;6:97–8.

© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Mahajan S, Ghosh GC, George OK. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-227257

Figure 1 Suprasternal view demonstrating vertical vein. AO, aorta; LA, left atrium; LPA, left pulmonary artery; MPA, main pulmonary artery; PV, pulmonary vein; RPA, right pulmonary artery; RSVC, right superior vena cava; VV, vertical vein.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow